



US006440424B1

(12) **United States Patent**
Sasaki et al.

(10) Patent No.: **US 6,440,424 B1**
(45) Date of Patent: ***Aug. 27, 2002**

(54) **HIGH MOLECULAR WEIGHT MAJOR
OUTER MEMBRANE PROTEIN OF
MORAXELLA**

(75) Inventors: **Ken Sasaki; Robin E. Harkness**, both
of Willowdale; **Sheena M. Loosmore**,
Aurora; **Michel H. Klein**, Willowdale,
all of (CA)

(73) Assignee: **Aventis Pasteur Limited**, Toronto (CA)

(*) Notice: This patent issued on a continued pro-
secution application filed under 37 CFR
1.53(d), and is subject to the twenty year
patent term provisions of 35 U.S.C.
154(a)(2).

Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **08/483,855**

(22) Filed: **Jun. 7, 1995**

Related U.S. Application Data

(63) Continuation-in-part of application No. 08/431,718, filed on
May 1, 1995, now Pat. No. 6,335,018.

(51) Int. Cl.⁷ **A61K 39/02; A61K 38/00;**
C07K 1/00; C07K 14/00

(52) U.S. Cl. **424/251.1; 530/350; 530/412;**
530/413; 530/414; 530/415; 424/190.1;
424/184.1; 424/234.1; 424/251.1; 514/2;
514/8; 930/200

(58) Field of Search **530/350, 412,**
530/413, 414, 415; 424/190.1, 251.1, 184.1,
234.1; 514/2, 8; 930/200

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,258,029 A	3/1981	Moloney et al.	424/49
4,855,283 A	8/1989	Lockhoff et al.	514/8
5,552,146 A	9/1996	Hansen	424/251.1
5,759,813 A *	6/1998	Hansen et al.	424/184.1
6,121,427 A *	9/2000	Huang et al.	435/320.1
6,214,981 B1 *	4/2001	Tucker et al.	
6,310,190 B1 *	10/2001	Hansen et al.	536/23.1
6,335,018 B1 *	1/2002	Sasaki et al.	424/184.1

FOREIGN PATENT DOCUMENTS

WO	WO A 93 03761	3/1883
WO	WO A 91 09952	7/1991
WO	WO A 93 10214	5/1993

OTHER PUBLICATIONS

Ostle et al. Am J. Vet. Res 47(7): 1419-1421, 1986 (Jul.).
Van Hare, G.F., P.A. Shurin, C.D. Marchant, N.A. Cartelli,
C.E. Johnson, D. Fulton, S. Carlin, and C.H. Kim. Acute
otitis media caused by *Branhamella catarrhalis*: biology and
therapy. Rev. Infect. Dis. 9:16-27.

Chapman, A.J., D.M. Musher, S. Jonsson, J.E. Clarridge,
and R.J. Wallace. 1985. Development of bactericidal anti-
body during *Branhamella catarrhalis* infection. J. Infect.
Dis. 151:878-882.

Hager, H., A. Verghese, S. Alvarez, and S.L. Berk. 1987.
Branhamella catarrhalis respiratory infections. Rev. Infect.
Dis. 9:1140-1149.

McLeod, D.T., F. Ahmad, M.J. Croughan, and M.A. Calder.
1986. Bronchopulmonary infection due to *M. catarrhalis*.
Clinical features and therapeutic response. Drugs
31(Suppl.3): 109-112.

Nicotra, B., M. Rivera, J.I. Luman, and R.J. Wallace. 1986.
Branhamella catarrhalis as a lower respiratory tract patho-
gen in patients with chronic lung disease. Arch.Intern.Med.
146:890-893.

Ninane, G., J. Joly, and M. Kraytman. 1978. Bronchopul-
monary infection due to *Branhamella catarrhalis* 11 cases
assessed by transtracheal puncture. Br.Med.Jr. 1:276-278.

Srinivasan, G., M.J. Raff, W.C. Templeton, S.J. Givens, R.C.
Graves, and J.C. Mel. 1981. *Branhamella catarrhalis* pneu-
monia. Report of two cases and review of the literature.
Am.Rev. Respir. Dis. 123:553-555.

West, M., S.L. Berk, and J.K. Smith. 1982. *Branhamella*
catarrhalis pneumonia. South.Med. J. 75:1021-1023.

Brorson, J-E., A. Axelsson, and S.E. Holm. 1976. Studies on
Branhamella catarrhalis (*Neisseria catarrhalis*) with spe-
cial reference to maxillary sinusitis. Scan. J. Infect. Dis.
8:151-155.

Evans, F.O., Jr., J.B. Sydnor, W.E.C. Moore, G.R. Moore,
J.L. Manwaring, A.H. Brill, R.T. Jackson, S. Hanna, J.S.
Skaar, L.V. Holdeman, G.S. Fitz-Hugh, M.A. Sande, and
J.M. Gwaltney, Jr. 1975. Sinusitis of the maxillary antrum.
N.Engl.J.Med. 293:735-739.

Tinkelman, D.G., and H.J. Silk. 1989. Clinical and bacte-
riologic features of chronic sinusitis in children. Am.J.Dis.
Child. 143:938-942.

Wald, E.R., C. Byers, N. Guerra, M.Casselbrant, and D.
Beste. 1989. Subacute sinusitis in children. J.Pediatr.
115:28-32.

(List continued on next page.)

Primary Examiner—Nita Minnifield

(74) Attorney, Agent, or Firm—Sim & McBurney

(57) **ABSTRACT**

An isolated and purified outer membrane protein of a
Moraxella strain, particularly *M. catarrhalis*, has a molecu-
lar mass of about 200 kDa. The about 200 kDa outer
membrane protein as well as nucleic acid molecules encod-
ing the same are useful in diagnostic applications and
immunogenic compositions, particularly for in vivo admin-
istration to a host to confer protection against disease caused
by a bacterial pathogen that produces the about 200 kDa
outer membrane protein or produces a protein capable of
inducing antibodies in a host specifically reactive with the
about 200 kDa outer membrane protein.

13 Claims, 17 Drawing Sheets

OTHER PUBLICATIONS

- Wald, E.R., G.J. Milmo, A. Bowen, J. Ledesma-Medina, N. Salamon, and C.D. Bluestone. 1981. Acute maxillary sinusitis in children. *N.Engl.J.Med.* 304:749-754.
- Christensen, J.J., and B. Bruun. 1985. Bacteremia caused by a beta-lactamase producing strain of *Branhamella catarrhalis*. *Acta. Pathol. Microbiol. Immunol. Scand. Sect. B.* 93:273-275.
- Craig, D.B., and P.A. Wehrle. 1983. *Branhamella catarrhalis* septic arthritis. *J. Rheumatol.* 10:985-986.
- Gray, L.D., R.E. Van Scoy, J.P. Anhalt, and P.K.W. Yu. 1989. Wound infection caused by *Branhamella catarrhalis*. *J. Clin. Microbiol.* 27:818-820.
- Guthrie, R., K. Bakenhaster, R. Nelson, and R. Woskobnick. 1988. *Branhamella catarrhalis* sepsis: a case report and review of the literature. *J. Infect. Dis.* 158:907-908.
- Hiroshi, S., E.J. Anaissie, N. Khardori, and G.P. Bodey. 1988. *Branhamella catarrhalis* septicemia in patients with leukemia. *Cancer* 61:2315-2317.
- O'Neill, J.H., and P.W. Mathieson. 1987. Meningitis due to *Branhamella catarrhalis*. *Aust. N.Z. J. Med.* 17:241-242.
- Murphy, T.F. 1989. The surface of *Branhamella catarrhalis*: a systematic approach to the surface antigens of an emerging pathogen. *Pediatr. Infect. Dis. J.* 8:S75-S77.
- Klingman, K.L., and T.F. Murphy. 1994. Purification and characterization of a high-molecular-weight outer membrane protein of *Moraxella (Branhamella) catarrhalis*. *Infect. Immun.* 62:1150-1155.
- Helminen, M.E., I. Maciver, J.L. Latimer, J. Klesney-Tait, L.D. Cope, M. Paris, G.H. McCracken, Jr., and E.J. Hansen. 1994. A large, antigenically conserved protein on the surface of *Moraxella catarrhalis* is a target for protective antibodies. *J. Infect. Dis.* 170:867-872.
- Panezutti H., O. James, E.J. Hanson, Y. Choi, R.E. Harkness, M.H. Klein and P. Chong, 1993. Identification of surface-exposed B-cell epitopes recognized by *Haemophilus influenzae* type b P1 specific monoclonal antibodies. *Infect. Immun.* 61: 1867-1872.
- Nixon-George et al. (1990), *J. Immunology* 144:4798-4802.
- Wiesmuller (1989), *Vaccine* 8:29-33.
- Deres et al. (1989), *Nature* 342:561.
- Lockhoff, O. Glycolipids as Immunomodulators: Synthesis and Properties. 1991. *Chem. Int. Ed. Engl.* 30:1611-1620.
- Journal of Infectious Diseases*, 158 (4). 1988. 761-765., XP002013102 Bartos L C et al: "Comparison of the Outer Membrane Proteins of 50 Strains of *Branhamella-catarrahilis*" see the whole document.
- Science, Apr. 14 1995, 268 (5208) P221-5, United States, XP002013103 Casey PJ: "Protein lipidation in cell signaling," see the whole document.
- Murphy et al; *Microbial Path.* 1989, 6:159-174.
- Burgess et al. *J. Cell Biology* 111:2129-38.
- Lazar et al. *Mol Cell Biology*, vol. 8 No. 3: 1247-52.
- Ostle et al., *AM J Vet Res*, vol. 47 No. 7: 1419-21.

* cited by examiner

Figure 1A




Figure 1B




Figure 2





Figure 3


Figure 4

FIG. 5.




FIG. 6A

Nucleotide Sequence Between SalI and NcoI

10	20	30	40	50	60	70
CCATGGATAT	GGCAGGTGT	GCTCGCCTGC	CGTATGATGG	CGATGACACC	CCATTGGCCC	CATATCTGTA
80	90	100	110	120	130	140
CGATTTGACA	TGTGATATGA	TTTAAACATGT	GACATGATTT	AACATGTGTT	AATACTGTTG	CCATCATTTAC
150	160	170	180	190	200	210
CATAAATTTAG	TAACGCATTT	AGTAACGCAT	TTGTAAAAAT	CATTGCGCCC	CTTTATGTGT	ATCATATGAA
220	230	240	250	260	270	280
TAGAATATTA	TGATTGTATC	TGATTATTGT	ATCAGAATGG	TGATGCTATA	TGATGATGCC	TACGAGTTGA
290	300	310	320	330	340	350
TTTGGGTTAA	TCACTCTATG	ATTTGATATA	TTTTGAAACT	AATCTATTGA	CTTAAATCAC	CATATGGTTA
360	370	380	390	400	410	420
TAATTTAGCA	TAATGGTAGG	CTTTTGTGTA	AAATCACATC	GCAATATTGT	TCTACTGTTA	CTACCATGCT
430	440	450	460	470	480	490
TGAATGACGA	TCCCAATCAC	CAGATTTCAT	CAAGTGATGT	GTTTGTATAC	GCACCATTTA	CCCTAATTAT
500	510	520	530	540	550	560
TTCAATCAAA	TGCCATATGC	AGCATGTATC	ATTTTTTTTAA	GGTAAACCAC	CATGAATCAC	ATCTATAAAG
570	580	590	600	610	620	630
TCATCTTTAA	CAAAGCCACA	GGCACATTTA	TGGCAGTGGC	AGAGTACGCC	AAATCCCACA	GCACGGGGGG

FIG. 6B

640	650	660	670	680	690	700
GGGGTAGCTG	TGCTACAGG	CAAGTTGGCA	GTGTATGCAC	TCTGAGCTTT	GCCCGTATTG	CCGCGCTCGC
710	720	730	740	750	760	770
TGTCCTCGTG	ATCGGTGCAA	CGCTCAGTGG	CAGTGCTTAT	GCTCAAAAAA	AAGATAACCA	ACATATCGCA
780	790	800	810	820	830	840
ATTGGTGAAC	AAAACCAGCC	AAGACGCTCA	GGCACTGCCA	AGGCGGACGG	TGATCGAGCC	ATTGCTATTG
850	860	870	880	890	900	910
GTGAAAATGC	TAACGCACAG	GGCGGTCAAG	CCATCGCCAT	CGGTAGTAGT	AATAAACTG	TCAATGGAAG
920	930	940	950	960	970	980
CAGTTTGGAT	AAGATAGGTA	CCGATGCTAC	GGGTCAAGAG	TCCATCGCCA	TCGGTGGTGA	TGTAAAGGCT
990	1000	1010	1020	1030	1040	1050
AGTGGTGATG	CCTCGATTGC	CATCGGTAGT	GATGACTTAC	ATTTGCTTGA	TCAGCATGGT	AATCCTAAAC
1060	1070	1080	1090	1100	1110	1120
ATCCGAAAGG	TACTCTGATT	AACGATCTTA	TTAACGGCCA	TGCAGTATTA	AAAGAAATAC	GAAGCTCAAA
1130	1140	1150	1160	1170	1180	1190
GGATAATGAT	GTAAAAATATA	GACGCACAAC	CGCAAGCGGA	CACGCCAGTA	CTGCAGTGGG	AGCCATGTCA
1200	1210	1220	1230	1240	1250	1260
TATGCACAGG	GTCAATTTTC	CAACGCCCTT	GGTACACGGG	CAACAGCTAA	AAGTGCCTAT	TCCTTGGCAG

FIG. 6C

1270	TGGGTC	1280	CGCCACAGCC	1290	GAGGCCAAAT	1300	CTACAATCGC	1310	TATTGGTTCT	1320	GATGCAACAT	1330	CTAGCTCGTT
1340	GGGAGCGGATA	1350	GCCCTTGGTG	1360	CAGGTA	1370	TGCTCAGCTA	1380	CAGGGCAGTA	1390	TTGCCCTAGG	1400	TCAGGTCT
1410	GTTGTCACTC	1420	AGAGTGATAA	1430	TAATTTCTAGA	1440	CCGGCCCTATA	1450	CACCAAAATAC	1460	CCAGGCACTA	1470	GACCCCAAGT
1480	TTCAAGCCAC	1490	CAATAATACG	1500	AAGCGGGTC	1510	CACCTTTCCAT	1520	TGGTAGTAAC	1530	TCTATCAAAC	1540	GTAAAAATCAT
1550	CAATGTCGGT	1560	GCAGGTGTTA	1570	ATAAAACCGA	1580	TGCGGTCAAT	1590	GTGGCACACG	1600	TAGAAGCGGT	1610	GGTGAAGTGG
1620	GCTAAGGAGC	1630	GTAGAATTAC	1640	TTTTTCAGGGT	1650	GATGATAACA	1660	GTACTGACGT	1670	AAAAATAGGT	1680	TTGGATAATA
1690	CCTTTAACTAT	1700	TAAAGGTGGT	1710	GCAGAGACCA	1720	ACGCATTAAAC	1730	CGATAATAAT	1740	ATCGGTGTGG	1750	TAAAAGAGGC
1760	TGATAATAGT	1770	GGTCTGAAAG	1780	TTAAACTTGC	1790	TAAAACTTTA	1800	AACAATCTTA	1810	CTGAGGTGAA	1820	TACAACTACA
1830	TTAAATGCCA	1840	CAACCACAGT	1850	TAAGGTAGGT	1860	AGTAGTAGTA	1870	GTACTACAGC	1880	TGAATTATTG	1890	AGTGATAGTT

FIG. 6D

1900	TAACCTTTAC	1910	CCAGCCCAAT	1920	ACAGGCAGTC	1930	AAAGCACAAG	1940	CAAAACCGTC	1950	TATGGCGTTA	1960	ATGGGGTGAA
1970	GTTTACTAAT	1980	AATGCAGAAA	1990	CAACAGCAGC	2000	AATCGGCACT	2010	ACTCGTATTA	2020	CCAGAGATAA	2030	AATTGGCTTT
2040	GCTCAGATG	2050	GTGATGTTGA	2060	TGAAAACAA	2070	GCACCATATT	2080	TGGATAAAA	2090	ACAACCTTAA	2100	GTGGGTAGTG
2110	TTGCAATTAC	2120	CATAGACAAT	2130	GGCATTGATG	2140	CAGGTAATAA	2150	AAAGATCAGT	2160	AATCTTGCCA	2170	AAGGTAGCAG
2180	TGCTAACGAT	2190	GCGGTTACCA	2200	TCGAACAGCT	2210	CAAAGCCGCC	2220	AAGCCTACTT	2230	TAAACGCAGG	2240	CGCTGGCATC
2250	AGTGTCACAC	2260	CTACTGAAAT	2270	ATCAGTTGAT	2280	GCTAAGAGTG	2290	GCAATGTTAC	2300	CGCCCCAACT	2310	TACAACATTG
2320	GCGTGAAAC	2330	CACCGAGCTT	2340	AACAGTGATG	2350	GCACTAGTGA	2360	TAAATTTAGT	2370	GTTAAGGGTA	2380	GTGGTACGAA
2390	CAATAGCTTA	2400	GTTACCGCCG	2410	AACATTGGC	2420	AAGCTATCTA	2430	AATGAAGTCA	2440	ATCGAACGGC	2450	TGACAGTGCT
2460	CTACAAAGCT	2470	TTACCGTTAA	2480	AGAAGAAGAC	2490	GATGATGACG	2500	CCAACGCTAT	2510	CACCGTGGCT	2520	AAAGATACGA

FIG. 6E

2530	2540	2550	2560	2570	2580	2590
CAAAAAATGC	CGGCGCAGTC	AGCATCTTAA	AACTCAAAGG	TAAAAACGGT	CTAACGGTTG	CTACCAAAAA
2600	2610	2620	2630	2640	2650	2660
AGATGGGTACG	GTTACCTTTG	GGCTTAGCCA	AGATAGCGGT	CTGACCATTG	GCAAAAGCAC	CCTAAACAAC
2670	2680	2690	2700	2710	2720	2730
GATGGCTTGA	CTGTAAAGA	TACCAACGAA	CAAATCCAAG	TCGGTGCTAA	TGGCATTAAA	TTTACTAATG
2740	2750	2760	2770	2780	2790	2800
TGAATGGTAG	TAATCCAGGT	ACTGGCATTG	CAAATACCGC	TCGCATTACC	AGAGATAAAA	TTGGCTTTGC
2810	2820	2830	2840	2850	2860	2870
TGGTTCTGAT	GGTGCAGTTG	ATACAAACAA	ACCTTATCTT	GATCAAGACA	AGCTACAAGT	TGGCAATGTT
2880	2890	2900	2910	2920	2930	2940
AAGATTACCA	ACACTGGCAT	TAACGCAGGT	GGTAAAGCCA	TCACAGGGCT	GTCCCCAACA	CTGCCTAGCA
2950	2960	2970	2980	2990	3000	3010
TTGCCGATCA	AAGTAGCCGC	AACATAGAAC	TGGGCAATAC	AATCCAAGAC	AAAGACAAAT	CCAACGCTGC
3020	3030	3040	3050	3060	3070	3080
CAGCATTAAT	GATATATTAA	ATACAGGCTT	TAAACCTAAAA	AATAATAACA	ACCCCATTTGA	CTTTGTCTCC
3090	3100	3110	3120	3130	3140	3150
ACTTATGACA	TTGTTGACTT	TGCCAATGGC	AATGCCACCA	CCGCCACAGT	AACCCATGAT	ACCGCTAACA

FIG. 6F

3160	3170	3180	3190	3200	3210	3220
AAACCAGTAA	AGTGGTATAT	GATGTGAATG	TGGATGATAC	AACCATTCAT	CTAACAGGCA	CTGATGACAA
3230	3240	3250	3260	3270	3280	3290
TAAAAAACTT	GGCGTCAAAA	CCACCAAACT	GAACAAAAA	AGTGCTAATG	GTAATACAGC	AACATACTTT
3300	3310	3320	3330	3340	3350	3360
AATGTTAACT	CTAGTGATGA	AGATGCCCTT	GTTAACGCCA	AAGACATCGC	CGAAAAATCTA	AACACCCCTAG
3370	3380	3390	3400	3410	3420	3430
CCAAGGAAAT	TCACACCCACC	AAAGGCACAG	CAGACACCGC	CCTACAAACC	TTTACCGTTA	AAAAGGTAGA
3440	3450	3460	3470	3480	3490	3500
TGAAAAATAAT	AATGCTGATG	ACGCCAACGC	CATCACCCGTG	GGTCAAAAAG	ACGCCAAATAA	TCAAGTCAAC
3510	3520	3530	3540	3550	3560	3570
ACCCTAACAC	TCAAAGGTGA	AAACGGTCTT	AATATTAAAA	CCGACAAAAA	TGGTACGGTT	ACCTTTGGCA
3580	3590	3600	3610	3620	3630	3640
TTAACACCCAC	AAGCGTCTT	AAAGCCGGCA	AAAGCACCCCT	AAACGACGGT	GGCTTGCTTA	TTAAAAAACCC
3650	3660	3670	3680	3690	3700	3710
CACTGGTAGC	GAACAAATCC	AAGTCGGTGC	TGATGGCGTG	AAGTTTGCCA	AGGTTAATAA	TAATGGTGTT
3720	3730	3740	3750	3760	3770	3780
GTAGGTGCTG	GCATTGATGG	CACAACCTCG	ATTACCAGAG	ATGAAATTGG	CTTTACTGGG	ACTAATGGCT

FIG. 6G

3790	3800	3810	3820	3830	3840	3850
CACCTTGATAA	AAGCAAACCC	CACCTAAGCA	AAGACGGCAT	TAACGCAGGT	GGTAAAAAGA	TTACCAACAT
3860	3870	3880	3890	3900	3910	3920
TCAATCAGGT	GAGATTGCC	AAAACAGCCA	TGATGCTGTG	ACAGGGCGCA	AGATTATGA	TTTTAAAAACC
3930	3940	3950	3960	3970	3980	3990
GAACTTGAAA	ACAAAATCAG	CAGTACTGCC	AAAACAGCAC	AAAACCTCAT	ACACGAATTC	TCAGTAGCAG
4000	4010	4020	4030	4040	4050	4060
ATGAACAAGG	TAATAACTTT	ACGGTTAGTA	ACCCTTACTC	CAGTTATGAC	ACCTCAAGA	CCTCTGATGT
4070	4080	4090	4100	4110	4120	4130
CATCACCTTT	GCAGGTGAAA	ACGGCATTTAC	CACCAAGGTA	AATAAAGGTG	TGGTGCGTGT	GGGCATTGAC
4140	4150	4160	4170	4180	4190	4200
CAAAACCAAG	GCTTAACCCAC	GCCTAAGCTG	ACCGTGGGTA	ATAATAATGG	CAAAGGCATT	GTCATTGACA
4210	4220	4230	4240	4250	4260	4270
GCCAAAATGG	TCAAAAATACC	ATCACAGGAC	TAAGCAACAC	TCTAGCTAAT	GTTACCAATG	ATAAAGGTAG
4280	4290	4300	4310	4320	4330	4340
CGTACGCACC	ACAGAACAGG	GCAATATAAT	CAAAGACGAA	GACAAAACCC	GTGCCGCCAG	CATTGTTGAT
4350	4360	4370	4380	4390	4400	4410
GTGCTAAGCG	CAGGCTTTAA	CTTGCAAGGC	AATGGTGAAG	CGGTTGACTT	TGCTCTCCACT	TATGACACCG

FIG. 6H

4420	TCAACTTTGC	4430	CGATGGCAAT	4440	GCCACCACCG	4450	CTAAGTGAC	4460	CTATGATGAC	4470	ACAAGCAAAA	4480	CCAGTAAAGT
4490	GGTCTATGAT	4500	GTC AATGTGG	4510	ATGATACAAC	4520	CATTGAAGTT	4530	AAAGATAAAA	4540	AAC TTGGCGT	4550	AAAAACCACC
4560	ACATTGACCA	4570	G TACTGGCAC	4580	AGGTGCTAAT	4590	AAATTTGCCC	4600	TAAGCAATCA	4610	AGCTACTGGC	4620	GATGCGGCTTG
4630	TCAAGGCCAG	4640	TGATATCGTT	4650	GCTCATCTAA	4660	ACACCTTATC	4670	TGGCGACATC	4680	CAAACTGCCA	4690	AAGGGGCAAG
4700	CCAAGCGAAC	4710	A ACTCAGCAG	4720	GCTATGTGGA	4730	TGCTGATGGC	4740	AATAAGGTCA	4750	TCTATGACAG	4760	TACCGATAAC
4770	AAGTACTATC	4780	AAGCCAAAAA	4790	TGATGGCACA	4800	GTTGATAAAA	4810	CCAAAGAAGT	4820	TGCCAAAGAC	4830	AAACTGGTCCG
4840	CCCAAGCCCA	4850	AACCCAGAT	4860	GGCACATTGG	4870	CTCAAAATGAA	4880	TGTCAAATCA	4890	GTCATTAACA	4900	AAGAACCAAGT
4910	AAATGATGCC	4920	AATAAAAAGC	4930	AAGGCATCAA	4940	TGAAGACAAC	4950	GCCTTTGTTA	4960	AAGGACTTGA	4970	AAAAGCCCGCT
4980	TCTGATAACA	4990	AAACCAAAAA	5000	CGCCGCAGTA	5010	ACTGTGGGTG	5020	ATTTAAATGC	5030	CGTTGCCCAA	5040	ACACCCGCTGA

FIG. 6I

5050	5060	5070	5080	5090	5100	5110
CCTTTGCAGG	GGATACAGC	ACAAAGGCTA	AAAACTGGG	CGAGACTTTG	ACCATCAAAG	GTGGGCAAAC
5120	5130	5140	5150	5160	5170	5180
AGACACCAAT	AAGCTAACCG	ATAATAACAT	CGGTGTGGTA	GCAGGTACTG	ATGGCTTCAC	TGTCAAACCTT
5190	5200	5210	5220	5230	5240	5250
GCCAAAGACC	TAACCAATCT	TAACAGCGTT	AATGCAGGTG	GCACCAAAAT	TGATGACAAA	GGCGTGTCTT
5260	5270	5280	5290	5300	5310	5320
TTGTAGACTC	AAGCGGTCAA	GCCAAAGCAA	ACACCCCTGT	GCTAAGTGCC	AATGGGCTGG	ACCTGGGTGG
5330	5340	5350	5360	5370	5380	5390
CAAGGTCATC	AGTAATGTGG	GCAAAGGCAC	AAAAGATACC	GACGCTGCCA	ATGTACAACA	GTTAACGAA
5400	5410	5420	5430	5440	5450	5460
GTACGCAACT	TGTTGGGTCT	TGGTAATGCT	GGTAATGATA	ACGCTGACGG	CAATCAGGTA	AACATTGCCG
5470	5480	5490	5500	5510	5520	5530
ACATCAAAA	AGACCCAAAT	TCAGGTTTCAT	CATCTAACCG	CACTGTTCATC	AAAGCAGGCA	CGGTACTTGG
5540	5550	5560	5570	5580	5590	5600
CGGTAAAGGT	AATAACGATA	CCGAAAAACT	TGCCACTGGT	GGTATACAAG	TGGGCGTGGG	TAAAGACGGC
5610	5620	5630	5640	5650	5660	5670
AACGCTAACG	GCGATTTAAG	CAATGTTTGG	GTCAAAAACCC	AAAAGATGG	CAGCAAAAAA	GCCCTGCTCG
5680	5690	5700	5710	5720	5730	5740
CCACTTATAA	CGCCGCAGGT	CAGACCAACT	ATTTGACCAA	CAACCCCGCA	GAAGCCATTG	ACAGAATAAA


FIG. 6J

5750	5760	5770	5780	5790	5800	5810
TGAACAAGGT	ATCCGCTTCT	TCCATGTCAA	CGATGGCAAT	CAAGAGCCTG	TGGTACAAGG	GGCTAACGGC
5820	5830	5840	5850	5860	5870	5880
ATTGACTCAA	GTGCCCTCAGG	CAAGCACTCA	GTGGCGGATAG	GTTTCCAGGC	CAAGGCAGAT	GGTGAAGCCG
5890	5900	5910	5920	5930	5940	5950
CCGTGCCAT	AGGCAGACAA	ACCCAAGCAG	GCAACCAATC	CATCGCCATC	GGTGATAACG	CACAAGCCAC
5960	5970	5980	5990	6000	6010	6020
GGGCGATCAA	TCCATCGCCA	TCGGTACAGG	CAATGTGGTA	GCAGGTAAGC	ACTCTGGTGC	CATCGGCGAC
6030	6040	6050	6060	6070	6080	6090
CCAAGCACTG	TTAAGGCTGA	TAACAGTTAC	AGTGTGGGTA	ATAACAACCA	GTTTACCGAT	GCCACTCAAA
6100	6110	6120	6130	6140	6150	6160
CCGATGTCTT	TGGTGTGGGC	AATAACATCA	CCGTGACCGA	AAGTAACTCG	GTTGCCCCTTAG	GTTCAAACTC
6170	6180	6190	6200	6210	6220	6230
TGCCATCAGT	GCAGGCACAC	ACGCAGGCAC	ACAAGCCAAA	AAATCTGACG	GCACAGCAGG	TACAACCACC
6240	6250	6260	6270	6280	6290	6300
ACAGCAGGTG	CAACCCGTAC	GGTTAAAGGC	TTTGCTGGAC	AAACGGCGGT	TGGTGCGGTC	TCCGTGGGTG
6310	6320	6330	6340	6350	6360	6370
CCTCAGGTGC	TGAACGCCGT	ATCCAAAATG	TGGCAGCAGG	TGAGGTCAGT	GCACCCAGCA	CCGATGCGGT
6380	6390	6400	6410	6420	6430	6440
CAATGGTAGC	CAGTTGTACA	AAGCCACCCA	AAGCAATTGCC	AACGCAACCA	ATGAGCTTGA	CCATCGTATC

FIG. 6K

6450	6460	6470	6480	6490	6500	6510
CACCAAAACG	AAAATAAGGC	CAATGCAGGG	ATTTCATCAG	CGATGGCGAT	GGCGTCCATG	CCACAAGCCT
6520	6530	6540	6550	6560	6570	6580
ACATTCCCTGG	CAGATCCATG	GTTACCGGGG	GTATTGCCAC	CCACAACGGT	CAAGGTGCGG	TGGCAGTGGG
6590	6600	6610	6620	6630	6640	6650
ACTGTCGAAG	CTGTCGGATA	ATGGTCAATG	GGTATTAA	ATCAATGGTT	CAGCCGATAC	CCAAGGCCAT
6660	6670	6680	6690	6700	6710	6720
GTAGGGGCGG	CAGTTGGTGC	AGGTTTTTCAC	TTTTAAGCCA	TAAATCGCAA	GATTTTACTT	AAAAATCAAT
6730	6740	6750	6760	6770	6780	6790
CTCACCATAG	TTGTATAAAA	CAGCATCAGC	ATCAGTCATA	TTACTGATGC	TGATGTTTTT	TATCACTTAA
6800	6810	6820	6830	6840	6850	6860
ACCATTTTAC	CGCTCAAGTG	ATTCTCTTTC	ACCATGACCA	AATCGCCATT	GATCATAGGT	AAACTTATTG
6870	6880	6890	6900	6910	6920	6930
AGTAAATTTT	ATCAATGTAG	TTGTTAGATA	TGGTTAAAT	TGTGCCATTG	ACCAAAAAAT	GACCGATTTA
6940	6950	6960	6970			
TCCCCGAAAT	TTCTGATTAT	GATCCGTTGA	CCTGCAGGTC	GAC		

FIG. 7.



1

HIGH MOLECULAR WEIGHT MAJOR OUTER MEMBRANE PROTEIN OF MORAXELLA

REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of U.S. patent application Ser. No. 08/431,718 filed May 1, 1995 now U.S. Pat. No. 6,335,018.

FIELD OF THE INVENTION

The present invention relates to the field of immunology and is particularly concerned with outer membrane proteins from *Moraxella*, methods of production thereof, genes encoding such proteins and uses thereof.

BACKGROUND OF THE INVENTION

Otitis media is the most common illness of early childhood with approximately 70% of all children suffering at least one bout of otitis media before the age of seven. Chronic otitis media can lead to hearing, speech and cognitive impairment in children. It is caused by bacterial infection with *Streptococcus pneumoniae* (approximately 50%), non-typable *Haemophilus influenzae* (approximately 30%) and *Moraxella* (Branhamella) *catarrhalis* (approximately 20%). In the United States alone, treatment of otitis media costs between one and two billion dollars per year for antibiotics and surgical procedures, such as tonsillectomies, adenoidectomies and insertion of tympanostomy tubes. Because otitis media occurs at a time in life when language skills are developing at a rapid pace, developmental disabilities specifically related to learning and auditory perception have been documented in youngsters with frequent otitis media.

M. catarrhalis mainly colonizes the respiratory tract and is predominantly a mucosal pathogen. Studies using cultures of middle ear fluid obtained by tympanocentesis have shown that *M. catarrhalis* causes approximately 20% of cases of otitis media (ref. 1—Throughout this application, various references are referred to in parenthesis to more fully describe the state of the art to which this invention pertains. Full bibliographic information for each citation is found at the end of the specification, immediately preceding the claims. The disclosures of these references are hereby incorporated by reference into the present disclosure).

The incidence of otitis media caused by *M. catarrhalis* is increasing. As ways of preventing otitis media caused by pneumococcus and nontypeable *H. influenzae* are developed, the relative importance of *M. catarrhalis* as a cause of otitis media can be expected to further increase.

M. catarrhalis is also an important cause of lower respiratory tract infections in adults, particularly in the setting of chronic bronchitis and emphysema (refs. 2, 3, 4, 5, 6, 7, and 8). *M. catarrhalis* also causes sinusitis in children and adults (refs. 9, 10, 11, 12, and 13) and occasionally causes invasive disease (refs. 14, 15, 16, 17, 18, and 19).

Like other Gram-negative bacteria, the outer membrane of *M. catarrhalis* consists of phospholipids, lipopolysaccharide (LPS), and outer membrane proteins (OMPs). Eight of the *M. catarrhalis* OMPs have been identified as major components. These are designated by letters A through H, beginning with OMP A which has a molecular mass of 98 kDa to OMP H which has a molecular mass of 21 kDa (ref. 20).

Recently, a high-molecular-weight outer membrane protein of *M. catarrhalis* was purified and characterized (ref.

2

21). The apparent molecular mass of this protein varies from 350 kDa to 720 kDa as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). This protein appears to be an oligomer of much smaller proteins or subunits thereof of molecular mass 120 to 140 kDa and is antigenically conserved among strains of *Moraxella*.

A similar protein named UspA was also reported to be present on the surface of this species of bacteria with an apparent molecular mass of 300 to 400 kDa (ref. 22). Judging from the molecular mass, these two proteins may be the same.

M. catarrhalis infection may lead to serious disease. It would be advantageous to provide other outer membrane proteins for *M. catarrhalis* and genes encoding such proteins for use as antigens in immunogenic preparations including vaccines; carriers for other antigens and immunogens and the generation of diagnostic reagents.

SUMMARY OF THE INVENTION

The present invention is directed towards the provision of purified and isolated major outer membrane protein of *Moraxella catarrhalis* and other *Moraxella* strains, having an apparent molecular mass of about 200 kDa.

In accordance with one aspect of the invention, there is provided an isolated and purified, outer membrane protein of a *Moraxella* strain having a molecular weight of about 200 kDa, as determined by SDS-PAGE, or a fragment or an analog thereof. The outer membrane protein may be substantially in its native conformation (so as to have substantially retained the characteristic immunogenicity of the outer membrane protein in the *Moraxella* strain) and may be isolated from a *M. catarrhalis* strain, such as from *M. catarrhalis* 4223. Such isolated and purified about 200 kDa outer membrane protein is substantially free from non-200 kDa outer membrane protein, phospholipids and lipopolysaccharide of *Moraxella*. The about 200 kDa outer membrane protein is at least about 70 wt % pure, preferably at least about 90 wt % pure, and may be in the form of an aqueous solution thereof.

The present invention also provides a purified and isolated nucleic acid molecule encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, or a fragment or an analog of the outer membrane protein. The protein encoded by the nucleic acid molecule may encode a protein containing the amino acid sequence NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2) for *Moraxella catarrhalis* strain 4223 or containing the corresponding amino acid sequence from other *Moraxella* strains.

In a further aspect of the present invention, there is provided a purified and isolated nucleic acid molecule having a sequence selected from the group consisting of (a) the sequence set out in FIG. 6 (SEQ ID No: 1), or the complementary sequence thereto; (b) a sequence encoding an about 200 kDa protein of a strain of *Moraxella* and containing the amino acid sequence NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2), or the complementary sequence thereto; and (c) a nucleotide sequence which hybridizes under stringent conditions to any one of the sequences defined in (a) or (b). The nucleic acid preferably defined in (c) has at least about 90% sequence identity with any one of the sequences defined in (a) or (b).

The nucleic acid molecules provided herein may be included in a vector adapted for transformation of a host.

The nucleic acid molecules provided herein also may be included in an expression vector adapted for transformation of a host along with expression means operatively coupled to the nucleic acid molecule for expression by the host of the about 200 kDa outer membrane protein of a strain of *Moraxella* or the fragment or the analog of the outer membrane protein. A transformed host containing the expression vector is included within the invention, along with a recombinant outer membrane protein or fragment or analog thereof producible by the transformed host.

The expression means may include a nucleic acid portion encoding a leader sequence for secretion from the host of the outer membrane protein or the fragment or the analog of the outer membrane protein. The expression means may include a nucleic acid portion encoding a lipidation signal for expression from the host of a lipidated form of the outer membrane protein or the fragment or analog thereof.

The present invention further includes a live vector for delivery of the outer membrane protein of the invention or a fragment or analog thereof, comprising a vector containing the nucleic acid molecule provided herein. The live vector may be selected from the group consisting of *E. coli*, *Salmonella*, BCG, adenovirus, poxvirus, vaccinia and poliovirus.

In accordance with a further aspect of the present invention, there is provided a peptide having no less than six amino acids and no more than 150 amino acids and containing an amino acid sequence corresponding to a portion only of the outer membrane protein of the invention, or a fragment or analog thereof. The peptide may be one having the amino acid sequence NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2) for the *Moraxella catarrhalis* 4283 strain or the amino acid sequence for the corresponding peptide for other strains of *Moraxella*.

The present invention also provides an immunogenic composition comprising an immunoeffective amount of an active component, which may be the outer membrane protein or fragment or analog thereof, nucleic acid molecules, recombinant outer membrane proteins, fragments or analogs thereof, live vectors, and/or peptides as provided herein, along with a pharmaceutically acceptable carrier therefor with the active component producing an immune response when administered to a host, which may be a primate, particularly a human. The immunogenic composition may be formulated as a vaccine for in vivo administration to a host to confer protection against diseases caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein. In particular, the bacterial pathogen is a strain of *Moraxella*, particularly *M. catarrhalis*. The immunogenic composition may be formulated as a microparticle capsule, ISCOM or liposome preparation. The immunogenic composition may be used in combination with a targeting molecule for delivery to specific cells of the immune system as to mucosal surfaces. The immunogenic compositions of the invention (including vaccines) may further comprise at least one other immunogenic or immunostimulating material and the immunostimulating material may be at least one adjuvant. Suitable adjuvants for use in the present invention include, (but are not limited to) aluminum phosphate, aluminum hydroxide, QS21, Quil A, calcium phosphate, calcium hydroxide, zinc hydroxide, a glycolipid analog, and octadecyl ester of an amino acid, a muramyl dipeptide and a lipoprotein. Advantageous combinations of adjuvants are described in copending U.S. patent

application Ser. No. 261,194 filed Jun. 16, 1994, assigned to the assignee hereof and the disclosure of which is incorporated herein by reference. The invention further includes an antibody specific for the outer membrane protein provided herein producible by immunizing a host with an immunogenic composition provided herein.

In a further aspect of the invention, there is provided a method of generating an immune response in a host comprising administering thereto an immuno-effective amount of the immunogenic composition as provided herein. The immune response may be a humoral or a cell-mediated immune response. The immune response may provide protection to the host against diseases caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein. Hosts in which protection against disease may be conferred include primates including humans.

The present invention provides, in an additional aspect thereof, a method of producing a vaccine comprising administering the immunogenic composition provided herein to a test host to determine an amount and a frequency of administration of the active component to confer protection against disease caused by a bacterial pathogen that produces the about 200 kDa outer membrane protein or produces a protein capable of inducing antibodies in the host specifically reactive with the about 200 kDa outer membrane protein, and formulating the active component in a form suitable for administration to a treated host in accordance with said determined amount and frequency of administration. The treated host may be a human.

A further aspect of the present invention provides a method of determining the presence of nucleic acid encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, in a sample, comprising the steps of:

- (a) contacting the sample with the nucleic acid molecule provided herein to produce duplexes comprising the nucleic acid molecules and any said nucleic acid molecule encoding the outer membrane protein present in the sample and specifically hybridizable therewith; and
- (b) determining the production of the duplexes.

In yet a further aspect of the invention, there is provided a method of determining the presence of antibodies specifically reactive with outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample, comprising the steps of:

- (a) contacting the sample with the outer membrane protein as provided herein to produce complexes comprising the outer membrane protein and any said antibodies present in the sample specifically reactive therewith; and
- (b) determining production of the complexes.

In a further aspect of the invention, there is also provided a method of determining the presence of an outer membrane protein if a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample comprising the steps of:

- (a) immunizing a subject with the immunogenic composition as provided herein, to produce antibodies specific for the outer membrane protein;
- (b) contacting the sample with the antibodies to produce complexes comprising any outer membrane protein present in the sample and said outer membrane protein specific antibodies; and
- (c) determining production of the complexes.

5

The outer membrane protein may be part of a *Moraxella catarrhalis* strain.

The present invention provides, in a yet further aspect, a diagnostic kit for determining the presence of nucleic acid encoding an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, in a sample, comprising:

- (a) the nucleic acid molecule as provided herein;
- (b) means for contacting the nucleic acid with the sample to produce duplexes comprising the nucleic acid molecule and any said nucleic acid present in the sample and hybridizable with the nucleic acid molecule; and
- (c) means for determining production of the duplexes.

In yet a further aspect of the invention, there is provided a diagnostic kit for determining the presence of antibodies in a sample specifically reactive with the outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, comprising:

- (a) the outer membrane protein as provided herein;
- (b) means for contacting the outer membrane protein with the sample to produce complexes comprising the outer membrane protein and any said antibodies present in the sample; and
- (c) means for determining production of the complexes.

The invention also provides a diagnostic kit for detecting the presence of an outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, in a sample, comprising:

- (a) an antibody specific for the about 200 kDa outer membrane protein as provided herein;
- (b) means for contacting the antibody with the sample to produce a complex comprising the outer membrane protein and outer membrane-specific antibody; and
- (c) means for determining production of the complex.

In a further aspect of the invention, there is provided a method of producing an isolated and purified outer membrane protein of a strain of *Moraxella* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, comprising the steps of:

- (a) providing a cell mass of the *Moraxella* strain;
- (b) disrupting the cell mass to provide a cell lysate;
- (c) fractionating the cell lysate to provide a fraction containing the outer membrane protein substantially free from other cell lysate components, and
- (d) recovering said outer membrane protein.

The bacterial strain may be *Moraxella catarrhalis*. The cell lysate may be fractionated by gel electrophoresis.

In this application, the term "about 200 kDa protein" is used to define a family of outer membrane proteins of *M. catarrhalis* having molecular mass of between about 160 and 230 kDa and includes proteins having variations in their amino acid sequences including those naturally occurring in various strains of *Moraxella*. The purified and isolated DNA molecules comprising a gene having an open reading frame of the about 200 kDa protein of the present invention also include those encoding functional analogs of the about 200 kDa protein. In this application, a first protein or peptide is a "functional analog" of a second protein if the first protein is immunologically related to and/or has the same function as the second protein or peptide. The functional analog may be, for example, a fragment of the protein or a substitution, addition or deletion mutant thereof.

Advantages of the present invention include:

- a method for isolating purified about 200 kDa outer membrane protein of a *Moraxella* strain that produces the outer membrane protein, including *Moraxella catarrhalis*;

6

an isolated and purified about 200 kDa outer membrane protein isolatable from a *Moraxella* strain; and diagnostic kits and immunological reagents for specific identification of *Moraxella* and hosts infected thereby.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A and 1B show an analysis of *Moraxella catarrhalis* cell proteins by SDS-PAGE. The identification of the lanes and the sources of the proteins are given in Example 2 below;

FIG. 2 shows a comparative analysis of cell proteins from a number of *M. catarrhalis* strains by SDS-PAGE analysis. The identification of the lanes and the sources of the proteins are given in Example 4 below;

FIG. 3 shows an analysis of isolated and purified about 200 kDa outer membrane protein of *M. catarrhalis* by SDS-PAGE. The identification of the lanes is given in Example 4 below;

FIG. 4 shows the specific recognition of about 200 kDa outer membrane protein by anti-peptide antiserum. The identification of the lanes and antiserum are given in Example 8 below;

FIG. 5 shows restriction maps of clones containing a gene having an open reading frame of the about 200 kDa outer membrane protein of *M. catarrhalis*. An open reading frame of the about 200 kDa outer membrane protein is indicated by the hatched box;

FIG. 6 shows the nucleotide sequence (SEQ ID No: 1) of the gene having an open reading frame of the about 200 kDa outer membrane protein of *M. catarrhalis*;

FIG. 7 shows the specific identification of *M. catarrhalis* expressing the about 200 kDa outer membrane protein by guinea pig anti-200 kDa specific antiserum in contrast to other bacteria. Identification of the lanes and bacteria appears below.

GENERAL DESCRIPTION OF INVENTION

Referring to FIGS. 1A and 1B and FIG. 2, there is illustrated the separation of a novel outer membrane protein from a variety of strains of *M. catarrhalis* having a molecular mass about 200 kDa. FIG. 3 shows the isolated and purified outer membrane protein.

The purified protein was eluted from the gel and used to raise antibodies in guinea pigs. The antibodies specifically recognize only strains of *M. catarrhalis* which produce the outer membrane protein (Table I below).

Referring to FIG. 4, there is shown the recognition of the about 200 kDa outer membrane protein by antibodies raised in guinea pigs to a synthesized peptide corresponding to an internal fragment of the about 200 kDa protein. The synthesized peptide had the amino acid sequence NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No: 3).

Referring to FIG. 5, there is shown restriction maps of clones containing a gene having an open reading frame of the about 200 kDa outer membrane protein. The nucleotide sequence (SEQ ID No: 1) of the gene having an open reading frame coding for the about 200 kDa outer membrane protein as shown in FIG. 6.

In one embodiment of the present invention, the isolated and purified about 200 kDa outer membrane protein as provided herein is useful for generating antibodies that can be used to specifically distinguish *M. catarrhalis* from other bacterial pathogens that cause otitis media and other dis-

7

eases. Thus referring to FIG. 7, there is illustrated an immunoblot showing the specific reactivity of a guinea pig monospecific anti-200 kDa outer membrane protein antiserum produced by immunizing mice with the purified about 200 kDa outer membrane protein as provided herein. The bacterial lysates analyzed were as follows:

Lane	Bacterium	Source
1.	Molecular Weight Standard	
2.	<i>M. catarrhalis</i> 4223	middle ear fluid
3.	<i>M. catarrhalis</i> RH408	non-clumping variant of strain 4223
4.	<i>H. influenzae</i> , MinnA strain	meningitis isolate
5.	non-typable <i>H. influenzae</i> , SB12 strain	otitis media isolate
6.	non-typable <i>H. influenzae</i> , SB33 strain	otitis media isolate
7.	<i>S. pneumoniae</i> type 6	ATCC 6306
8.	<i>S. pneumoniae</i> type 14	ATCC 6314
9.	<i>P. aeruginosa</i>	
10.	<i>E. coli</i> DH5a	

The results shown in FIG. 7 clearly show the usefulness of outer membrane-specific antisera as provided herein to distinguish between bacterial pathogens that produce diseases with similar clinical symptoms.

Thus, in accordance with another aspect of the present invention, there is provided a vaccine against *Moraxella*, comprising an immunogenically-effective amount of the outer membrane protein as provided herein and a physiologically-acceptable carrier therefor. The outer membrane protein provided herein also may be used as a carrier protein for hapten, polysaccharides or peptides to make a conjugate vaccine against antigenic determinants unrelated to the about 200 kDa outer membrane protein.

The about 200 kDa outer membrane protein provided herein is useful as a diagnostic reagent, as an antigen for the generation of anti-outer membrane protein antibodies, or as an antigen for vaccination against the diseases caused by species of *Moraxella* for detecting infection by *Moraxella*.

In additional embodiments of the present invention, the about 200 kDa outer membrane protein as provided herein may be used as a carrier molecule to prepare chimeric molecules and conjugate vaccines (including glycoconjugates) against pathogenic bacteria, including encapsulated bacteria. Thus, for example, glycoconjugates of the present invention may be used to confer protection against disease and infection caused by any bacteria having polysaccharide antigens including lipooligosaccharides (LOS) and polyribosylphosphate (PRP). Such bacterial pathogens may include, for example, *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Escherichia coli*, *Neisseria meningitidis*, *Salmonella typhi*, *Streptococcus mutants*, *Cryptococcus neoformans*, *Klebsiella*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Particular antigens which can be conjugated to outer membrane protein and methods to achieve such conjugations are described in published PCT application WO 94/12641, assigned to the assignee hereof and the disclosure of which is hereby incorporated by reference thereto.

In another embodiment, the carrier function of the outer membrane protein may be used, for example, to induce immunity toward abnormal polysaccharides of tumor cells, or to produce anti-tumor antibodies that can be conjugated to chemotherapeutic or bioactive agents.

It is clearly apparent to one skilled in the art, that the various embodiments of the present invention have many applications in the fields of vaccination, diagnosis, treatment

8

of *Moraxella* infections, and in the generation of immunological reagents. A further non-limiting discussion of such uses is further presented below.

1. Vaccine Preparation and Use

Immunogenic compositions, suitable to be used as vaccines, may be prepared from the about 200 kDa outer membrane protein as disclosed herein, which may be purified from the bacteria or which may be produced recombinantly, as well as immunological fragments thereof. The vaccine elicits an immune response in a subject which produces antibodies, including anti-200 kDa outer membrane protein antibodies and antibodies that are opsonizing or bactericidal. Should the vaccinated subject be challenged by *Moraxella* or other bacteria that produce proteins capable of producing antibodies that specifically recognize 200 kDa outer membrane protein, the antibodies bind to and inactivate the bacterium. Furthermore, opsonizing or bactericidal anti-200 kDa outer membrane protein antibodies may also provide protection by alternative mechanisms.

Immunogenic compositions including vaccines may be prepared as injectables, as liquid solutions or emulsions. The about 200 kDa outer membrane protein may be mixed with pharmaceutically acceptable excipients which are compatible with the about 200 kDa outer membrane protein. Such excipients may include, water, saline, dextrose, glycerol, ethanol, and combinations thereof. The immunogenic compositions and vaccines may further contain auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, or adjuvants to enhance the effectiveness thereof. Immunogenic compositions and vaccines may be administered parenterally, by injection subcutaneously or intramuscularly. Alternatively, the immunogenic compositions formed according to the present invention, may be formulated and delivered in a manner to evoke an immune response at mucosal surfaces. Thus, the immunogenic composition may be administered to mucosal surfaces by, for example, the nasal or oral (intra-gastric) routes. Alternatively, other modes of administration including suppositories and oral formulations may be desirable. For suppositories, binders and carriers may include, for example, polyalkylene glycols or triglycerides. Oral formulations may include normally employed excipients such as, for example, pharmaceutical grades of saccharine, cellulose and magnesium carbonate. These compositions can take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 1 to 95% of the about 200 kDa outer membrane protein. The immunogenic preparations and vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective, protective and immunogenic. The quantity to be administered depends on the subject to be treated, including, for example, the capacity of the individual's immune system to synthesize antibodies, and if needed, to produce a cell-mediated immune response. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are readily determinable by one skilled in the art and may be of the order of micrograms of the about 200 kDa outer membrane protein. Suitable regimes for initial administration and booster doses are also variable, but may include an initial administration followed by subsequent administrations. The dosage may also depend on the route of administration and will vary according to the size of the host.

The concentration of the about 200 kDa outer membrane antigen in an immunogenic composition according to the

invention is in general about 1 to 95%. A vaccine which contains antigenic material of only one pathogen is a monovalent vaccine. Vaccines which contain antigenic material of several pathogens are combined vaccines and also belong to the present invention. Such combined vaccines contain, for example, material from various pathogens or from various strains of the same pathogen, or from combinations of various pathogens.

Immunogenicity can be significantly improved if the antigens are co-administered with adjuvants, commonly used as 0.05 to 0.1 percent solution in phosphate-buffered saline. Adjuvants enhance the immunogenicity of an antigen but are not necessarily immunogenic themselves. Adjuvants may act by retaining the antigen locally near the site of administration to produce a depot effect facilitating a slow, sustained release of antigen to cells of the immune system. Adjuvants can also attract cells of the immune system to an antigen depot and stimulate such cells to elicit immune responses.

Immunostimulatory agents or adjuvants have been used for many years to improve the host immune responses to, for example, vaccines. Intrinsic adjuvants, such as lipopolysaccharides, normally are the components of the killed or attenuated bacteria used as vaccines. Extrinsic adjuvants are immunomodulators which are typically non-covalently linked to antigens and are formulated to enhance the host immune responses. Thus, adjuvants have been identified that enhance the immune response to antigens delivered parenterally. Some of these adjuvants are toxic, however, and can cause undesirable side-effects, making them unsuitable for use in humans and many animals. Indeed, only aluminum hydroxide and aluminum phosphate (collectively commonly referred to as alum) are routinely used as adjuvants in human and veterinary vaccines. The efficacy of alum in increasing antibody responses to diphtheria and tetanus toxoids is well established and, more recently, a HBsAg vaccine has been adjuvanted with alum. While the usefulness of alum is well established for some applications, it has limitations. For example, alum is ineffective for influenza vaccination and inconsistently elicits a cell mediated immune response.

A wide range of extrinsic adjuvants can provoke potent immune responses to antigens. These include saponins complexed to membrane protein antigens (immune stimulating complexes), pluronic polymers with mineral oil, killed mycobacteria in mineral oil, Freund's complete adjuvant, bacterial products, such as muramyl dipeptide (MDP) and lipopolysaccharide (LPS), as well as lipid A, and liposomes.

To efficiently induce humoral immune responses (HIR) and cell-mediated immunity (CMI), immunogens are emulsified in adjuvants. Many adjuvants are toxic, inducing granulomas, acute and chronic inflammations (Freund's complete adjuvant) FCA, cytotoxicity (saponins and Pluronic polymers) and pyrogenicity, arthritis and anterior uveitis (LPS and MDP). Although FCA is an excellent adjuvant and widely used in research, it is not licensed for use in human or veterinary vaccines because of its toxicity.

Desirable characteristics of ideal adjuvants include:

- (1) lack of toxicity;
- (2) ability to stimulate a long-lasting immune response;
- (3) simplicity of manufacture and stability in long-term storage;
- (4) ability to elicit both CMI and HIR to antigens administered by various routes, if required;
- (5) synergy with other adjuvants;

(6) capability of selectively interacting with populations of antigen presenting cells (APC);

(7) ability to specifically elicit appropriate T_H1 or T_H2 cell-specific immune responses; and

(8) ability to selectively increase appropriate antibody isotype levels (for example, IgA) against antigens. U.S. Pat. No. 4,855,283 granted to Lockhoff et al on Aug. 8, 1989 which is incorporated herein by reference thereto, teaches glycolipid analogues including N-glycosylamides, N-glycosylureas and N-glycosylcarbamates, each of which is substituted in the sugar residue by an amino acid, as immunomodulators or adjuvants. Thus, Lockhoff et al. (U.S. Pat. No. 4,855,283 and ref. 27) reported that N-glycolipid analogs displaying structural similarities to the naturally-occurring glycolipids, such as glycosphingolipids and glycoacylglycerolipids, are capable of eliciting strong immune responses in both herpes simplex virus vaccine and pseudorabies virus vaccine. Some glycolipids have been synthesized from long chain-alkylamines and fatty acids that are linked directly with the sugars through the anomeric carbon atom, to mimic the functions of the naturally occurring lipid residues.

U.S. Pat. No. 4,258,029 granted to Moloney, assigned to the assignee hereof and incorporated herein by reference thereto, teaches that octadecyl tyrosine hydrochloride (OTH) functioned as an adjuvant when complexed with tetanus toxoid and formalin inactivated type I, II and III poliomyelitis virus vaccine. Also, Nixon-George et al. (ref. 24), reported that octadecyl esters of aromatic amino acids complexed with a recombinant hepatitis B surface antigen, enhanced the host immune responses against hepatitis B virus.

Lipidation of synthetic peptides has also been used to increase their immunogenicity. Thus, Wiesmuller (ref. 25) describes a peptide with a sequence homologous to a foot-and-mouth disease viral protein coupled to an adjuvant tripalmityl-S-glyceryl-cysteinylserine, being a synthetic analogue of the N-terminal part of the lipoprotein from Gram negative bacteria. Furthermore, Deres et al. (ref. 26) reported in vivo priming of virus-specific cytotoxic T lymphocytes with synthetic lipopeptide vaccine which comprised of modified synthetic peptides derived from influenza virus nucleoprotein by linkage to a lipopeptide, N-palmitoyl-S-[2,3-bis(palmitoylxy)-(2RS)-propyl-[R]-cysteine (TPC).

2. Immunoassays

The about 200 kDa outer membrane protein of the present invention is useful as an immunogen for the generation of anti-200 kDa outer membrane protein antibodies, as an antigen in immunoassays including enzyme-linked immunosorbent assays (ELISA), RIAs and other non-enzyme linked antibody binding assays or procedures known in the art for the detection of anti-bacterial, anti-Moraxella, and anti-200 kDa outer membrane protein antibodies. In ELISA assays, the about 200 kDa outer membrane protein is immobilized onto a selected surface, for example, a surface capable of binding proteins such as the wells of a polystyrene microtiter plate. After washing to remove incompletely adsorbed about 200 kDa outer membrane protein, a nonspecific protein such as a solution of bovine serum albumin (BSA) that is known to be antigenically neutral with regard to the test sample may be bound to the selected surface. This allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific bindings of antisera onto the surface.

The immobilizing surface is then contacted with a sample, such as clinical or biological materials, to be tested in a manner conducive to immune complex (antigen/antibody) formation. This may include diluting the sample with diluents, such as solutions of BSA, bovine gamma globulin (BGG) and/or phosphate buffered saline (PBS)/Tween. The sample is then allowed to incubate for from 2 to 4 hours, at temperatures such as of the order of about 20° to 37° C. Following incubation, the sample-contacted surface is washed to remove non-immunocomplexed material. The washing procedure may include washing with a solution, such as PBS/Tween or a borate buffer. Following formation of specific immunocomplexes between the test sample and the bound about 200 kDa outer membrane protein, and subsequent washing, the occurrence, and even amount, of immunocomplex formation may be determined by subjecting the immunocomplex to a second antibody having specificity for the first antibody. If the test sample is of human origin, the second antibody is an antibody having specificity for human immunoglobulins and in general IgG. To provide detecting means, the second antibody may have an associated activity such as an enzymatic activity that will generate, for example, a colour development upon incubating with an appropriate chromogenic substrate. Quantification may then be achieved by measuring the degree of colour generation using, for example, a visible spectrophotometer.

BIOLOGICAL MATERIALS

Certain plasmids that contain portions of the gene having the open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis* strain 4223 that are described and referred to herein have been deposited with the America Type Culture Collection (ATCC) located at 12301 Parklawn Drive, Rockville, Md., 20852, U.S.A., pursuant to the Budapest Treaty and prior to the filing of this application. The identification of the respective portions of the gene and their molecular size are contained in FIG. 5.

Samples of the deposited plasmids will become available to the public upon grant of a patent based upon this United States patent application. The invention described and claimed herein is not to be limited in scope by plasmids deposited, since the deposited embodiment is intended only as an illustration of the invention. Any equivalent or similar plasmids that encode similar or equivalent antigens as described in this application are within the scope of the invention.

Plasmid	ATCC Designation	Date Deposited
pKS47	97,111	April 7, 1995
pKS5	97,110	April 7, 1995
pKS9	97,114	April 18, 1995

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitations.

Methods of molecular genetics, protein biochemistry, and immunology used but not explicitly described in this dis-

closure and these Examples are amply reported in the scientific literature and are well within the ability of those skilled in the art.

EXAMPLE 1

This Example illustrates the generation of a non-clumping strain (RH408) of *M. catarrhalis*.

M. catarrhalis strain 4223, a clumping strain (a common property of *Moraxella* strains), was inoculated into several flasks containing 20 mL of brain heart infusion (BHI) broth, and the cultures were incubated with shaking (170 rpm) overnight at 37° C. Five mL of each overnight culture were transferred to five individual 1-mL tubes, and were left sitting undisturbed at room temperature for 3 to 8 hours, to allow bacteria to sediment. One hundred μ L of the cleared upper phase of each culture medium were used to inoculate 25 mL of BHI broth and cultures were incubated overnight at 37° C., as described above. This passaging was repeated six times, using 25 μ L of cleared medium to inoculate 25 mL of BHI for each overnight culture. Non-clumping bacterial cultures were identified by measuring the absorbency A_{578} at intervals over a 3 hour time period, in order to compare the sedimentation rates of the passaged strains to that of the original *M. catarrhalis* strain 4223 culture. Non-clumping mutants, including *M. catarrhalis* RH408, did not aggregate during the three hour time period. On BHI agar plates, strain RH408 had a colony morphology typical for all non-clumping strains. Strain RH408 was previously deposited at the ATCC under the Budapest Treaty on Dec. 13, 1994 with Accession No. 55637.

EXAMPLE 2

This Example illustrates the identification of the above 200 kDa outer membrane protein of *Moraxella catarrhalis*.

M. catarrhalis strains 4223, RH408, 5191, 8185, M2, M5, ATCC 25240, 3, 56, 135, 585 were grown in brain heart infusion (BHI) broth. The culture was incubated overnight with aeration at 37° C.

M. catarrhalis cells were sonicated and the total protein was determined using the BCA assay system (Pierce, Rockford, Ill.). Ten μ g of total protein were mixed with the SDS-PAGE sample buffer containing 0.3M Tris-HCl (pH 8.0), 50% glycerol, 10% SDS, 20% α -mercaptoethanol and 0.01% bromophenol blue, boiled for 5 minutes and loaded on each lane of SDS-PAGE gel (0.75 mm thick, 7.5% acrylamide). The gels were run at 200 V for 1 hour. Proteins were visualized by staining gels with a solution containing 0.13% Coomassie brilliant blue R, 10% acetic acid and 45% methanol. Excess stain was removed with a destaining solution of 5% ethanol and 7.5% acetic acid.

The various *Moraxella* proteins separated by this procedure are shown in FIGS. 1A and 1B. The *M. catarrhalis* strains tested were as follows:

Lane	Bacterial Strain	Source
FIG. 1A		
1.	Molecular Weight Standards	
2.	<i>E. coli</i>	
3.	No sample	
4.	<i>M. catarrhalis</i> 4223	middle ear fluid non-clumping variant of 4223
5.	<i>M. catarrhalis</i> RH408	

13

-continued

Lane	Bacterial Strain	Source
6.	<i>M. catarrhalis</i> 5191	middle ear fluid
7.	<i>M. catarrhalis</i> 8185	nasopharynx
8.	<i>M. catarrhalis</i> M2	sputum
9.	<i>M. catarrhalis</i> M5	sputum
10.	<i>M. catarrhalis</i> 25240	ATCC 25240

FIG. 1B

1.	<i>E. coli</i>	
2.	No sample	
3.	Molecular Weight Size Markers	
4.	<i>M. catarrhalis</i> 4223	middle ear fluid
5.	<i>M. catarrhalis</i> RH408	non-clumping variant of 4223
6.	<i>M. catarrhalis</i> 3	sputum
7.	<i>M. catarrhalis</i> 56	sputum
8.	<i>M. catarrhalis</i> 135	middle ear fluid
9.	<i>M. catarrhalis</i> 585	Blood

The about 200 kDa outer membrane protein was clearly seen in all otitis media strains (*M. catarrhalis* 4223, 5191, 135), one isolate from nasopharynx (8185), and in one isolate from sputum (M2). However, the about 200 kDa protein was not detected in three isolates from sputum (3, 56 and M5) and in one strain with unknown origin (ATCC 25240). A very narrow band was found in an isolate from blood of a bacteremia patient (585) and this band was recognized by an anti-200 kDa specific guinea pig serum on an immunoblot. Strain RH408 is a non-clumping spontaneous mutant isolated from strain 4223 (see Example 1) and was found to not express the about 200 kDa protein.

EXAMPLE 3

This Example illustrates the detection of antibodies specific for the about 200 kDa outer membrane protein in a serum obtained from a convalescent patient having recovered from otitis media due to *M. catarrhalis*.

After separation by SDS-PAGE, bacterial proteins were transferred from acrylamide gels to prepared PVDF (polyvinylidene-fluoride; Millipore) membranes at a constant voltage of 70 V for 1.5 h in a buffer system consisting of 3 g Tris, 14.4 g glycine and 200 ml methanol per liter at 4° C. Membranes with transferred proteins were blocked with Blocking Reagent (from Boehringer Mannheim) diluted in TBS (0.1M Tris, 0.15M NaCl) at room temperature for 30 min. Blots were exposed to convalescent antiserum diluted 1:500 in Blocking Reagent/TBS with 0.1% Tween 20 for 2 hours at room temperature. This patient had otitis media and the *M. catarrhalis* strain isolated from the patient's ear fluid was *M. catarrhalis* CJ7. Blots were then washed 2 times in Blocking Reagent/TBS with Tween at 15 min per wash. The reporter conjugate, horseradish peroxidase (HRP) conjugated to protein G, was diluted 1:4000 with Blocking Reagent/TBS with Tween and used to immerse the washed membranes for 30 min at room temperature. Blots were washed twice as above, followed by a TBS wash. Bound antibodies were detected using the Lumi-Glo (Kirkegaard and Perry) chemiluminescent detection system as described by the manufacturer. Treated blots were exposed to X-ray film. Antibodies were detected in this convalescent serum that reacted with the about 200 kDa outer membrane protein of *M. catarrhalis* CJ7. These results indicate that the about 200 kDa outer membrane protein is an immunogenic protein of *M. catarrhalis* to which an immune response is elicited during a natural infection by *M. catarrhalis*.

14

EXAMPLE 4

This Example illustrates the isolation and purification of the about 200 kDa outer membrane protein.

M. catarrhalis 4223 cells were harvested by centrifugation at 2,000 rpm for 10 min and frozen. The frozen cells were thawed, resuspended in 20 mM sodium phosphate buffer (pH 7.2) and sonicated until the cells were disrupted. The frozen-thawed cells were also lysed in 20 mM Tris buffer (pH 8) containing 4% SDS and 0.2 mM EDTA by boiling for 5 min to produce a cell lysate. The cell sonicates and cell lysates were suspended in a SDS-polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer, boiled for 5 min and separated by SDS-PAGE on a gel (1.5 mm thick, 7.5% acrylamide). The estimated position of about 200 kDa protein on the gel was cut out and the protein extracted from the gel by electroelution using the same buffer as the SDS-PAGE running buffer. The isolated about 200 kDa outer membrane protein was shown to be a homogeneous, single band by SDS-PAGE as seen in FIG. 3. The samples analyzed in FIG. 3 are as follows:

Lane	Sample
1.	Molecular Weight Size Markers
2.	Isolated and purified 200 kDa outer membrane protein

The isolated and purified 200 kDa outer membrane protein of *M. catarrhalis* shown in FIG. 3 has a purity of at least 70%. Purified about 200 kDa outer membrane protein preparations of at least 95% could be readily achieved.

When gels were run longer, they showed heterogeneity in the apparent molecular masses of the about 200 kDa outer membrane protein in different strains of *M. catarrhalis* (FIG. 2). In FIG. 2 the strains analyzed were as follows:

Lane	Strain	Source
1.	Molecular Weight Size Markers	
2.	<i>M. catarrhalis</i> H04	middle ear fluid
3.	<i>M. catarrhalis</i> H12	middle ear fluid
4.	<i>M. catarrhalis</i> PO34	middle ear fluid
5.	<i>M. catarrhalis</i> PO51	middle ear fluid
6.	<i>M. catarrhalis</i> E-07	middle ear fluid
7.	<i>M. catarrhalis</i> E-22	middle ear fluid
8.	<i>M. catarrhalis</i> E-23	middle ear fluid
9.	<i>M. catarrhalis</i> RH 4223	middle ear fluid
10.	<i>M. catarrhalis</i> RH 408	Non-clumping variant of 4223

The strain H12 (lane 3) was a natural isolate from middle ear fluid, but did not produce the about 200 kDa protein.

There may be at least three different sizes of protein in the about 200 kDa range. However, antibodies raised against the about 200 kDa outer membrane protein from one strain of *M. catarrhalis* (4223) did recognize all about 200 kDa proteins tested, present in different strains of *M. catarrhalis*. It is possible, however, that in particular immunogenic compositions, for example, as a vaccine and in particular diagnostic embodiments, that the about 200 kDa outer membrane protein from a variety of *M. catarrhalis* isolates (including immunogenically diverse isolates) may be required.

EXAMPLE 5

This Example illustrates the immunization of guinea pigs with purified about 200 kDa protein from *M. Catarrhalis*.

15

Approximately 30 to 40 μ g of the about 200 kDa protein, which was isolated from *M. catarrhalis* strain 4223 by electroelution, were mixed with Freund's complete adjuvant (FCA) and was subcutaneously injected into guinea pigs. After two weeks, the animals were boosted with about the same amount of the about 200 kDa protein in incomplete Freund's adjuvant (IFA). Two weeks later, blood was collected from the guinea pigs and antisera were obtained.

One antiserum was examined on Western blot for its reactivity with the about 200 kDa protein present in 54 different strains of *M. catarrhalis*, which were isolated in different geographical locations throughout the world (Canada, U.S. and Finland) (see Table 1 below). The about 200 kDa protein band was clearly recognized by the antiserum in all strains, in which the presence of the about 200 kDa protein band was detected on SDS-PAGE gels stained with Coomassie Blue. These results indicate that common epitopes of the about 200 kDa protein were conserved in all *M. catarrhalis* strains, which possessed this protein. As stated earlier, this protein is not present in all *M. catarrhalis* strains, but almost all strains, which were isolated from middle ear fluids from otitis media patients, did possess this protein (Table 1).

EXAMPLE 6

This Example illustrates the specific recognition of *M. catarrhalis* strain 4223 with anti-200 kDa protein guinea pig serum by ELISA assay (see Table 2 below).

M. catarrhalis strains 4223, RH408 (200 kDa negative mutant) and H-12 were cultured in 60 mL BHI broth overnight. *E. coli* strain BL21 (DE3) was cultured in 60 mL Luria-Bratani (LB) broth overnight. The cultures were split into three tubes and centrifuged. *M. catarrhalis* strain 4223 was centrifuged at 1,500 rpm for 10 minutes, H-12 at 2,000 rpm for 10 minutes, and RH408 and *E. coli* BL21 (DE3) at 3,000 rpm for 15 minutes. The pellet in one tube was suspended in 20 ml of Dulbecco's phosphate buffered saline (D-PBS) and diluted to $\frac{1}{500}$ with coating buffer (0.05M carbonate/bicarbonate buffer) pH 9.6. One hundred μ L of the bacteria solution was distributed in each well and incubated for 1 hour at room temperature. One hundred μ L of 0.2% glutaraldehyde was added to each well and incubated at room temperature for 10 minutes to fix the cells on the well. The wells were washed with PBS containing 0.1% Tween 20 and 0.1% BSA (washing buffer), and then blocked with PBS containing 0.1% BSA for 30 minutes at room temperature. After washing 5 times for 10 seconds with the washing buffer, serial dilutions of guinea pig antiserum with the washing buffer were added to wells and incubated at room temperature for 60 minutes. After washing, goat anti-guinea pig IgG conjugated with horseradish peroxidase was added to each well at the dilution of $\frac{1}{20,000}$. After incubation at room temperature for 60 minutes, the wells were washed and then color reaction was developed using 3,3',5,5'-tetramethylbenzidine (TMB) and hydrogen peroxide.

The ELISA plate wells were also coated with sonicates containing 10 μ g/mL of total proteins in the coating buffer, blocked without the fixation process and then assayed as described above.

The results shown in Table 2 indicate that the 200 kDa outer membrane protein specific guinea pig antiserum specifically recognizes strains of *M. catarrhalis* which produce the about 200 kDa protein. The ability of the antiserum to recognize whole cells indicates that the protein is present on the surface of the bacterial cells.

EXAMPLE 7

This Example describes the determination of an internal amino acid sequence of the 200 kDa outer membrane protein.

16

The about 200 kDa outer membrane protein was isolated from *M. catarrhalis* 4223 as described above. The protein was subjected to CNBr degradation, the proteolytic digests subjected to SDS-PAGE and transferred onto PVDF membrane. A peptide band migrating at a position corresponding to approximately 40 kDa was cut out from the membrane and its N-terminal amino acid sequence was determined. In another experiment, the CNBr degradation products of the about 200 kDa protein were subjected to a direct determination of N-terminal amino acid sequencing without separating by SDS-PAGE. Both analyses gave an identical, N-terminal sequence of 20 amino acids with one unidentified amino acid at the 17th position. The internal sequence of the 200 kDa outer membrane protein was:

NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-X-Gln-Gly-Ile (SEQ ID No: 2).

EXAMPLE 8

This Example describes the immunization of guinea pigs with a peptide corresponding to an internal fragment of the about 200 kDa outer membrane protein and the analysis of the antiserum generated.

Based upon the determination of the amino acid sequence of an internal fragment of the about 200 kDa outer membrane protein, a 16 amino acid long peptide of sequence:

NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No: 3)

was synthesized by standard procedure. This 16-mer peptide was conjugated to KLH using Imject Maleimide Activated KLH (Pierce, Rockford, Ill.) and approximately 500 μ g of the conjugate was injected into guinea pigs using the same immunization and boosting schedule as described above. The guinea pig serum raised against the 16-mer internal amino acid sequence (SEQ ID No: 3) was examined by immunoblot analysis and found to specifically recognize 200 kDa outer membrane protein in cell sonicates of *M. catarrhalis* 4223. The results are shown in FIG. 4 and indicate that the anti-peptide guinea pig antiserum specifically recognizes the 200 kDa protein of *M. catarrhalis* 4223. The samples analyzed in FIG. 4 were as follows:

Lane	Sample	Antiserum
1.	Molecular Weight Markers	
2.	Purified 200 kDa outer membrane protein	Anti-200 kDa protein
3.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:500
4.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:100
5.	<i>M. catarrhalis</i> cell sonicate	Anti-peptide 1:500
6.	<i>M. catarrhalis</i> cell sonicate	Pre-immune serum

The results obtained confirm that the peptide corresponding to SEQ ID Nos: 2 and 3 are derived from the 200 kDa outer membrane protein.

EXAMPLE 9

This Example describes the preparation of a *M. catarrhalis* library.

Chromosomal DNA was isolated as follows:

M. catarrhalis cell pellet was resuspended in 20 mL Tris-EDTA (TE) buffer, pH 7.5. Pronase (final concentration 500 μ g/mL) and SDS (final concentration 1%) were added and the suspension was incubated in a 37° C. water bath for 2 hours. DNA was isolated by sequential extractions with phenol (1 time), phenol-chloroform (1:1, 2 times), and

chloroform-isoamyl alcohol (24:1, 1 time). Extracted DNA was dialyzed against 1M NaCl at 4° C. for 4 hours. This was followed by dialysis against TE buffer, pH 7.5, at 4° C. for 48 hours (3 buffer changes). DNA was ethanol precipitated from the dialysate. Large size DNA was collected by spooling on a glass rod. Air dried DNA was dissolved in 3 mL water. Small scale Sau3A (New England BioLabs) restriction digests of chromosomal DNA (final volume 10 µl) were done to establish conditions required to obtain maximal amounts of chromosomal DNA with a size range of 15–23 kb. Large scale digests were prepared once the optimal digestion conditions were defined. The large scale digests consisted of 50 µl chromosomal DNA (290 µg/mL), 33 µl water, 10 µl Sau3A buffer (New England BioLabs), 1 µl BSA (10 mg/ml, New England BioLabs) and 6.3 µl Sau3A (0.04 U/µL), and were incubated at 37° C. for 15 minutes. Reactions were stopped with the addition of 10 µl 10xloading buffer (100 mM Tris-HCl pH 8, 10 mM EDTA, 0.1% bromophenol blue, 50% glycerol). Digested DNA was applied to 0.5% agarose gels (prepared in Tris-acetate-EDTA (TAE)) and separated according to size at 50 V for 6 hours. The region of the gel encompassing DNA of size 15–23 kb was cut from the gel and placed in dialysis tubing (BRL) with 3 mL TAE. DNA was electroeluted from the gel slice overnight at a field strength of 1 V/cm. Electroeluted DNA in TAE was extracted once with phenol, once with phenol-chloroform (1:1), and finally precipitated with ethanol. The dried DNA pellet was dissolved in 5 µL water. Size-fractionated chromosomal DNA was ligated with BamHI cut EMBL3 arms (Promega) using T4 DNA ligase in a final volume of 9 µL. The entire ligation reaction was packaged into phage λ using a commercially purchased packaging kit following the manufacturer's (Amersham) protocol.

The packaged DNA library was amplified on solid medium. This was accomplished by incubating 0.1 ml *E. coli* strain NM539 plating cells (cells suspended in 10 mM MgSO₄) with 15–25 µL of the packaged DNA library at 37° C. for 15 minutes. Bacteria with adsorbed phage were plated onto BBL plates (10 g BBL trypticase peptone, 5 g NaCl and 15 g agar per liter) using 3 mL of BBL top-agarose (same as BBL plates except agar replaced with 0.6% agarose) and plates were incubated overnight at 37° C. Phage were eluted from the top-agarose by adding 3 mL SM buffer (50 mM Tris-HCl, pH 7.5, 8 mM MgSO₄, 100 mM NaCl, 0.01% gelatin) to the plates and leaving them at 4° C. for 7 hours. SM buffer containing phage was collected from the plates, transferred to a screwcap tube and stored at 4° C. over chloroform.

EXAMPLE 10

This Example describes the cloning on a gene encoding the *M. catarrhalis* 200 kDa outer membrane protein.

The *M. catarrhalis* genomic library in phage lambda EMBL3 was screened using an anti-200 kDa protein guinea pig antiserum. A lambda phage clone 8II (FIG. 5), which expressed as about 200 kDa protein, was confirmed by immunoblotting of the phage lysate using the about 200 kDa outer membrane-specific antiserum.

Plate lysate cultures of this recombinant phage were prepared. The DNA was extracted from the plate lysates using a Wizard Lambda Preps DNA Purification System (Promega Corp., Madison, Wis.) according to the manufacturer's instructions. This phage clone carried a DNA insert of about 16 kb in size (the restriction map for which is in FIG. 5). The phage DNA was digested with a mixture of two restriction enzymes, Sall and XhoI, and separated by agar-

ose gel electrophoresis. Two DNA bands, approximately 5 kb and 11 kb in size, respectively, were cut out from the gel and extracted using a GeneClean kit (BIO 101 Inc., LaJolla, Calif.) according to a manufacturer's instruction.

The smaller 5 kb fragment was ligated into a plasmid vector, pBluescript II SK +/- (Stratagene Cloning Systems, LaJolla, Calif.), which was previously digested with Sall and XhoI, to produce plasmid pKS5. The larger 11 kb fragment was ligated into a plasmid vector, pSP72 (Promega Corp., Madison, Wis.), to produce plasmid pKS9.

Both ligated plasmids were used to transform *E. coli*, strain DH5α.

The lambda phage DNA was also digested with a mixture of XhoI and KpnI and the approximately 1.2 kb fragment was isolated after agarose gel separation as described above. This 1.2 kb fragment was ligated into a plasmid vector, pGEM-7Zf(+) (Promega Corp., Madison, Wis.), to produce plasmid pKS47. Restriction maps of the lambda and plasmid clones are shown in FIG. 5.

EXAMPLE 11

This Example describes the sequencing of the gene having an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis*.

The gene encoding the about 200 kDa outer membrane protein was sequenced by an Applied Biosystems sequencer. The one strand of the insert in the plasmid pKS5, was sequenced after construction of a nested set of deletions using a Erase-a-Base system (Promega Corp., Madison, Wis.). The plasmid pKS5 was first digested with XhoI and KpnI, treated with exonuclease III to generate a nested set of deletions in the insert and then recircularized according to the manufacturer's instructions. *E. coli* DH5α was transformed with a series of plasmids with deletions generated in this way. Plasmids were isolated from the transformants using a Quiagen midi plasmid isolation kit (Quiagen) and the size of plasmids examined by agarose gel electrophoresis after restriction enzyme digestion. The inserts of the plasmids with deletions were sequenced using a bacteriophage T7 promoter sequence as a primer.

Based upon the sequence, nucleotide primers were synthesized. Using the synthetic nucleotide primers, sequence gaps, which were not sequenced by the Erase-a Base system, were determined.

The sequence of the insert in plasmid pKS47 was determined from both ends using synthetic nucleotide primers. The nucleotide sequence of the gene has an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis* as shown in FIG. 6 (SEQ ID No: 1). This sequence included a nucleotide sequence:

5'-AATGTCAAATCAGTCATTAACAAAGAACAA-
GTAAATGATGCCAATAAAAAGCAAGGCATC-3'
(SEQ ID No: 4)

which encodes the internal amino acid sequence of the about 200 kDa outer membrane protein (SEQ ID No: 2) determined above. This result confirms that the cloned gene has an open reading frame of the gene coding for the about 200 kDa outer membrane protein of *M. catarrhalis*.

SUMMARY OF THE DISCLOSURE

In summary of the disclosure, the present invention provides an isolated and purified outer membrane protein of a Moraxella strain, particularly *M. catarrhalis*, having a molecular weight of about 200 kDa as well as isolated and

purified DNA molecules encoding the outer membrane protein. The invention also provides peptides corresponding to portions of the outer membrane protein. The protein, DNA sequences, recombinant proteins derived therefrom and peptides are useful for diagnosis, immunization and the generation of diagnostic and immunological reagents. Modifications are possible within the scope of this invention.

TABLE I

Presence of about 200 kDa outer membrane protein in various isolates of <i>Moraxella catarrhalis</i>		
Type of Clinical Isolate	Number of isolates Examined	Number of isolates ¹ containing the 200 kDa outer membrane protein
Otitis Media	37	36
Sputum/Expectoration/Bronchial Secretion	13	6
Blood	2	2
Nasopharynx	1	1
Unknown	1	0

¹The presence of the about 200 kDa outer membrane protein was determined by immunoblot analysis using a monospecific guinea pig anti-200 kDa protein antiserum.

TABLE II

Detection of about 200 kDa outer membrane protein of <i>M. catarrhalis</i> by the monospecific anti-200 kDa outer membrane guinea pig antiserum		
Strain	Sample	Reciprocal Reactive Titre
4223	Whole cells not fixed	800
RH408	Whole cells not fixed	<200
H12	Whole cells not fixed	<200
<i>E. coli</i> BL21	Whole cells not fixed	<200
4223		3200
RH408		200
H12		<200
<i>E. coli</i> BL21		<200
4223	Sonicate	12,800
RH408	Sonicate	800
H12	Sonicate	800
<i>E. coli</i> BL21	Sonicate	200

REFERENCES

- Van Hare, G. F., P. A. Shurin, C. D. Marchant, N. A. Cartelli, C. E. Johnson, D. Fulton, S. Carlin, and C. H. Kim. Acute otitis media caused by *Branhamella catarrhalis*: biology and therapy. *Rev. Infect. Dis.* 9:16-27.
- Chapman, A. J., D. M. Musher, S. Jonsson, J. E. Clarridge, and R. J. Wallace. 1985. Development of bactericidal antibody during *Branhamella catarrhalis* infection. *J. Infect. Dis.* 151:878-882.
- Hager, H., A. Verghese, S. Alvarez, and S. L. Berk. 1987. *Branhamella catarrhalis* respiratory infections. *Rev. Infect. Dis.* 9:1140-1149.
- McLeod, D. T., F. Ahmad, M. J. Croughan, and M. A. Calder. 1986. Bronchopulmonary infection due to *M. catarrhalis*. Clinical features and therapeutic response. *Drugs* 31(Suppl. 3):109-112.
- Nicotra, B., M. Rivera, J. I. Luman, and R. J. Wallace. 1986. *Branhamella catarrhalis* as a lower respiratory tract pathogen in patients with chronic lung disease. *Arch. Intern. Med.* 146:890-893.
- Ninane, G., J. Joly, and M. Kraytman. 1978. Bronchopulmonary infection due to *Branhamella catarrhalis* 11 cases assessed by transtracheal puncture. *Br. Med. J.* 1:276-278.
- Srinivasan, G., M. J. Raff, W. C. Templeton, S. J. Givens, R. C. Graves, and J. C. Mel. 1981. *Branhamella catarrhalis* pneumonia. Report of two cases and review of the literature. *Am. Rev. Respir. Dis.* 123:553-555.
- West, M., S. L. Berk, and J. K. Smith. 1982. *Branhamella catarrhalis* pneumonia. *South. Med. J.* 75:1021-1023.
- Brorson, J.-E., A. Axelsson, and S. E. Holm. 1976. Studies on *Branhamella catarrhalis* (*Neisseria catarrhalis*) with special reference to maxillary sinusitis. *Scan. J. Infect. Dis.* 8:151-155.
- Evans, F. O., Jr., J. B. Sydnor, W. E. C. Moore, G. R. Moore, J. L. Manwaring, A. H. Brill, R. T. Jackson, S. Hanna, J. S. Skaar, L. V. Holdeman, G. S. Fitz-Hugh, M. A. Sande, and J. M. Gwaltney, Jr. 1975. Sinusitis of the maxillary antrum. *N. Engl. J. Med.* 293:735-739.
- Tinkelman, D. G., and H. J. Silk. 1989. Clinical and bacteriologic features of chronic sinusitis in children. *Am. J. Dis. Child.* 143:938-942.
- Wald, E. R., C. Byers, N. Guerra, M. Casselbrant, and D. Beste. 1989. Subacute sinusitis in children. *J. Pediatr.* 115:28-32.
- Wald, E. R., G. J. Milmo, A. Bowen, J. Ledesma-Medina, N. Salamon, and C. D. Bluestone. 1981. Acute maxillary sinusitis in children. *N. Engl. J. Med.* 304:749-754.
- Christensen, J. J., and B. Bruun. 1985. Bacteremia caused by a beta-lactamase producing strain of *Branhamella catarrhalis*. *Acta. Pathol. Microbiol. Immunol. Scand. Sect. B* 93:273-275.
- Craig, D. B., and P. A. Wehrle. 1983. *Branhamella catarrhalis* septic arthritis. *J. Rheumatol.* 10:985-986.
- Gray, L. D., R. E. Van Scoy, J. P. Anhalt, and P. K. W. Yu. 1989. Wound infection caused by *Branhamella catarrhalis*. *J. Clin. Microbiol.* 27:818-820.
- Guthrie, R., K. Bakenhaster, R. Nelson, and R. Woskobnick. 1988. *Branhamella catarrhalis* sepsis: a case report and review of the literature. *J. Infect. Dis.* 158:907-908.
- Hiroshi, S., E. J. Anaissie, N. Khardori, and G. P. Bodey. 1988. *Branhamella catarrhalis* septicemia in patients with leukemia. *Cancer* 61:2315-2317.
- O'Neill, J. H., and P. W. Mathieson. 1987. Meningitis due to *Branhamella catarrhalis*. *Aust. N.Z. J. Med.* 17:241-242.
- Murphy, T. F. 1989. The surface of *Branhamella catarrhalis*: a systematic approach to the surface antigens of an emerging pathogen. *Pediatr. Infect. Dis. J.* 8: S75-S77.
- Klingman, K. L., and T. F. Murphy. 1994. Purification and characterization of a high-molecular-weight outer membrane protein of *Moraxella* (*Branhamella*) *catarrhalis*. *Infect. Immun.* 62:1150-1155.
- Helminen, M. E., I. Maciver, J. L. Latimer, J. Klesney-Tait, L. D. Cope, M. Paris, G. H. McCracken, Jr., and E. J. Hansen. 1994. A large, antigenically conserved protein on the surface of *Moraxella catarrhalis* is a target for protective antibodies. *J. Infect. Dis.* 170:867-872.
- Panzutti H., O. James, E. J. Hanson, Y. Choi, R. E. Harkness, M. H. Klein and P. Chong, 1993. Identification of surface-exposed B-cell epitopes recognized by *Haemophilus influenzae* type b P1 specific monoclonal antibodies. *Infect. Immun.* 61:1867-1872.
- Nixon-George et al. (1990), *J. Immunology* 144:4798-4802.
- Wiesmuller (1989), *Vaccine* 8:29-33.
- Deres et al. (1989), *Nature* 342:561.
- Lockhoff, O. Glycolipids as Immunomodulators: Synthesis and Properties. 1991. *Chem. Int. Ed. Engl.* 30:1611-1620.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 6973

<212> TYPE: DNA

<213> ORGANISM: Moraxella catarrhalis

<400> SEQUENCE: 1

```

ccatggatat gggcagggtg gctgcctgc cgtatgatgg cgatgacacc ccatttgccc      60
catatctgtg cgatttgaca tgtgatatga tttaacatgt gacatgattt aacattgttt      120
aatactgttg ccatacattac cataatttag taacgcattt agtaacgcat ttgtaaaaat      180
cattgcccgc ctttatgtgt atcatatgaa tagaattatta tgattgtatc tgattattgt      240
atcagaatgg tgatgctata tgatgatgcc tacgagttga tttgggttaa tcactctatg      300
atttgatata ttttgaaact aatctattga cttaaatcac catatgggta taatttagca      360
taatggtagg ctttttgtaa aaatcacatc gcaatattgt tctactgtta ctaccatgct      420
tgaatgacga tcccaatcac cagattcatt caagtgatgt gtttgatac gcaccattta      480
ccctaattat ttcaatcaaa tgcctatgtc agcatgtatc atttttttaa ggtaaacccac      540
catgaatcac atctataaag tcactcttaa caaagccaca ggcacattta tggcagtggc      600
agagtacgcc aaatcccaca gcacgggggg ggggtagctg tgctacaggg caagttggca      660
gtgtatgcac tctgagcttt gcccgatttg ccgcgctcgc tgctcctcgt atcggtgcaa      720
cgctcagtg g cagtgtctat gctcaaaaaa aagataccaa acatatacga attgggtgaa      780
aaaaccagcc aagacgtcca ggcactgcca agggggacgg tgatcgagcc attgctattg      840
gtgaaaatgc taacgcacag ggcggtcaag ccatacgccat cggtagtagt aataaaactg      900
tcaatggaag cagtttgatg aagataggtg ccgatgtac gggtaagag tccatcgcca      960
tcgggtggtg tgtaaaaggct agtggtagtg cctcgattgc catcggtagt gatgacttac      1020
atttgcttga tcagcatggt aatcctaacc atccgaaagg tactctgatt aacgatctta      1080
ttaacggcca tgcagtatta aaagaaatcc gaagctcaaa ggataatgat gtaaaatata      1140
gacgcacacg cgcaagcgga cacgccaagta ctgcagtggg agccatgtca tatgcacagg      1200
gtcatttttc caacgccttt ggtacacggg caacagctaa aagtgcctat tccctggcag      1260
tgggtcttgc cgccacagcc gagggccaat ctacaatcgc tattggttct gatgcaacat      1320
ctagctcgtt gggagcgata gcccttggtg caggtactcg tgctcagcta cagggcagta      1380
ttgccctagg tcaaggttct gttgtcactc agagtataa taattctaga ccggcctata      1440
caccataatc ccaggcacta gaccccaagt ttcaagccac caataatagc aaggcgggtc      1500
cactttccat tggtagtaac tctatcaaac gtaaaatcat caatgtcggg gcagggtgta      1560
ataaaaccga tgcggtcaat gtggcacagc tagaagcggg ggtgaagtgg gctaaggagc      1620
gtagaattac ttttcagggt gatgataaca gtactgacgt aaaaataggt ttggataata      1680
ctttaactat taaagggtgt gcagagacca acgcattaac cgataataat atcgtgtgtg      1740
taaaagaggg tgataaatgt ggtctgaaag ttaaaacttc taaaacttta aacaatctta      1800
ctgagggtgaa tacaactaca ttaaatgcca caaccacagt taaggtaggt agtagtagta      1860
gtactacagc tgaattattg agtgatagtt taacctttac ccagcccaat acaggcagtc      1920
aaagcacaa g caaaaccgtc tatggcggtt atgggggtgaa gtttactaat aatgcagaaa      1980
caacagcagc aatcggcact actcgtatta ccagagataa aattggcttt gctcgagatg      2040

```

-continued

gtgatgttga tgaanaacaa gcaccatatt tggataaaaa acaacttaaa gtgggtagt	2100
ttgcaattac catagacaat ggcattgatg caggtataaa aaagatcagt aatcttgcca	2160
aaggtagcag tgctaacgat gcggttacca tcgaacagct caaagccgcc aagcctactt	2220
taaacgcagg cgctggcacc agtgtcacac ctactgaaat atcagttgat gctaagagt	2280
gcaatgttac cgccccaaat tacaacattg gcgtgaaac caccgagctt aacagtgat	2340
gcactagtga taaatttagt gtttaagggtg gtggtacgaa caatagctta gttaccgccg	2400
aacatttggc aagctatcta aatgaagtca atcgaacggc tgacagtgtt ctacaaagct	2460
ttaccgttaa agaagaagac gatgatgacg ccaacgctat caccgtggct aaagatacga	2520
caaaaaatgc cggcgcagtc agcatcttaa aactcaagg taaaaacggt ctaacgggtg	2580
ctacaaaaaa agatggtagc gttactttg gcttagacca agatagcggt ctgaccattg	2640
gcaaaagcac cctaacaac gatggcttga ctgttaaaga taccaacgaa caaatccaag	2700
tcggtgctaa tggcattaaa tttactaatg tgaatggtag taatccaggt actggcattg	2760
caaataccgc tcgcattacc agagataaaa ttggctttgc tggttctgat ggtgcagttg	2820
atacaaaaa accttatctt gatcaagaca agctacaagt tggcaatgtt aagattacca	2880
acactggcat taacgcaggt ggtaaagcca tcacagggtt gtcccaaca ctgcctagca	2940
ttgccgatca aagtagccgc aacatagaac tgggcaatc aatccaagac aaagacaaat	3000
ccaacgctgc cagcattaat gatatttaa atacaggctt taacctaaaa aataataaca	3060
acccattga ctttgtctcc acttatgaca ttgttgactt tgccaatggc aatgccacca	3120
ccgccacagt aaccocatgat accgctaaca aaaccagtaa agtggatat gatgtgaatg	3180
tggatgatac aaccattcat ctaacaggca ctgatgacaa taaaaaactt ggcgtcaaaa	3240
ccacaaaact gaacaaaaca agtgctaag gtaatacagc aactaacttt aatgttaact	3300
ctagtgatga agatgcctt gttaacgcca aagacatcgc cgaatctta aacaccctag	3360
ccaaggaaat tcacaccacc aaaggcacag cagacaccgc cctacaaacc tttaccgtta	3420
aaaaggtaga tgaataaat aatgctgatg acgccaacgc catcacctgt ggtcaaaa	3480
acgcaataaa tcaagtcaac accctaacac tcaaagggtg aaacggtctt aatattaaaa	3540
ccgacaaaaa tggtagcgtt acctttggca ttaacaccac aagcgtctt aaagccggca	3600
aaagcacctt aaacgcaggt ggcttgtcta ttaaaaacc cactggtagc gaacaaatcc	3660
aagtcggtgc tgatggcgtg aagtttgcca aggttaataa taatgggtgt gtaggtgctg	3720
gcattgatgg cacaactcgc attaccagag atgaattgg cttactggg actaatggct	3780
cacttgataa aagcaaaccc cacctaagca aagacggcat taacgcaggt ggtaaaaaga	3840
ttaccaacat tcaatcaggt gagattgccc aaacagcca tgatgctgtg acaggcggca	3900
agatttatga tttaaaaacc gaacttgaaa acaaaatcag cagtactgcc aaacagcac	3960
aaaactcatt acacgaattc tcagtagcag atgaacagg taataacttt acggttagta	4020
accottactc cagttatgac acctcaaga cctctgatgt catcaccttt gcagggtgaa	4080
acggcattac caccaaggta aataaagggt tggtagcgtt gggcattgac caaaccaag	4140
gcttaaccac gcctaagctg accgtgggtg ataataatgg caaagcatt gtcattgaca	4200
gccaaaatgg tcaaaatacc atcacaggac taagcaacac tctagctaat gttaccaatg	4260
ataaaggtag cgtacgcacc acagaacagg gcaatataat caaagacgaa gacaaaacco	4320
gtgcccgcag cattgttgat gtgctaagcg caggctttaa cttgcaaggc aatggtgaa	4380
cggttgactt tgtctccact tatgacaccg tcaactttgc ctagggcaat gccaccaccg	4440

-continued

ctaagggtgac ctatgatgac acaagcaaaa ccagtaaagt ggtctatgat gtcaatgtgg	4500
atgatacaac cattgaagtt aaagataaaa aacttggcgt aaaaaccacc acattgacca	4560
gtactggcac aggtgctaataa aaatttggcc taagcaatca agctactggc gatgcgcttg	4620
tcaaggccag tgatatcggt gctcatctaa acaccttatac tggcgacata caaactgcca	4680
aaggggcaag ccaagcgaac aactcagcag gctatgtgga tgctgatggc aataaggtca	4740
tctatgacag taccgataac aagtactatc aagccaaaaa tgatggcaca gttgataaaa	4800
ccaangaagt tgccaagac aaactggctg cccaagccca aacccagat ggcacattgg	4860
ctcaaatgaa tgtcaaatca gtcattaaca aagaacaagt aaatgatgcc aataaaaagc	4920
aaggcatcaa tgaagacaac gcctttgtta aaggacttga aaaagccgct tctgataaca	4980
aaacaaaaaa cgccgcagta actgtgggtg atttaaatgc cgttgcacca acaccgctga	5040
cccttgacgg ggatacaggg acaacggcta aaaaactggg cgagactttg accatcaaa	5100
gtggggcaaac agacaccaat aagctaaccg ataataacat cgggtgtgta gcaggtactg	5160
atggcttcac tgtcaactt gccaaagacc taaccaatct taacagcgtt aatgcaggtg	5220
gcacaaaaat tgatgacaaa ggcgtgtctt ttgtagactc aagcggctca gccaaagcaa	5280
acacccctgt gctaagtgc aatgggctgg acctgggtgg caaggtcatc agtaatgtgg	5340
gcaaaggcac aaaagatacc gacgctgcca atgtacaaca gttaaagcaa gtacgcaact	5400
tggtgggtct tggtaatgct ggtaatgata acgctgacgg caatcaggta aacattgccg	5460
acatcaaaaa agacccaat tcaggttcat catctaaccg cactgtcatc aaagcaggca	5520
cggtaacttg cggtaaagg aataacgata ccgaaaaact tgccactggt ggtatacaag	5580
tggtcggtga taaagacggc aacgctaacy gcgatttaag caatgtttgg gtcanaacc	5640
aaaaagatgg cagcaaaaaa gccctgctcg ccacttataa cgccgcaggt cagaccaact	5700
atttgacca caaccccgca gaagccattg acagaataaa tgaacaaggt atccgcttct	5760
tccatgtcaa cgatggcaat caagagcctg tggtaacaag gcgtaacggc attgactcaa	5820
gtgcctcagg caagcactca gtggcgatag gtttccaggc caaggcagat ggtgaagccg	5880
ccgttgccat aggcagacaa acccaagcag gcaaccaatc catcgccatc ggtgataacg	5940
cacaagccac gggcgatcaa tccatcgcca tcggtacagg caatgtggtg gcaggtaaac	6000
actctggtgc catcgcgac ccaagcactg ttaaggctga taacagttac agtgtgggtg	6060
ataacaacca gtttaccgat gccactcaaa ccgatgtctt tgggtgtggc aataacatca	6120
ccgtgaccga aagtaactg tttgccttag gttcaaaact tgccatcagt gcaggcacac	6180
acgcaggcac acaagccaaa aaatctgacg gcacagcagg tacaaccacc acagcaggtg	6240
caaccggtac ggttaaggc tttgctggac aaacggcggt tgggtcggtc tccgtgggtg	6300
cctcaggtgc tgaacgccgt atccaaaatg tggcagcagg tgaggtcagt gccaccagca	6360
ccgatgcggt caatggtagc cagttgtaca aagccaccca aagcattgcc aacgcaacca	6420
atgagcttga ccactgtatc caccaaaacg aaataaaggc caatgcaggg atttcatcag	6480
cgatggcgat ggcgtccatg ccacaagcct acattcctgg cagatccatg gttaccgggg	6540
gtattgccac ccacaacggt caagggtcgg tggcagtggt actgtogaag ctgtcgata	6600
atggtcaatg ggtattttaa atcaatggt cagccgatac ccaaggccat gtggggcg	6660
cagttgtgc aggtttttac ttttaagcca taaatcgcaa gattttactt aaaaatcaat	6720
ctcaccatag ttgtataaaa cagcatcagc atcagtcata ttactgatgc tgatgtttt	6780
tatcacttaa accattttac cgtcaagt attctctttc accatgacca aatcgccatt	6840

-continued

gatcataggt aaacttattg agtaaatatt atcaatgtag ttgttagata tggtaaanaa 6900
 tgtgccattg accaaaaaat gaccgattta tcccgaanaa ttctgattat gatccgttga 6960
 cctgcaggtc gac 6973

<210> SEQ ID NO 2
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: *Moraxella catarrhalis*
 <220> FEATURE:
 <221> NAME/KEY: UNSURE
 <222> LOCATION: (17)

<400> SEQUENCE: 2

Asn Val Lys Ser Val Ile Asn Lys Glu Gln Val Asn Asp Ala Asn Lys
 1 5 10 15

Xaa Gln Gly Ile
 20

<210> SEQ ID NO 3
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: *Moraxella catarrhalis*

<400> SEQUENCE: 3

Asn Val Lys Ser Val Ile Asn Lys Glu Gln Val Asn Asp Ala Asn Lys
 1 5 10 15

<210> SEQ ID NO 4
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: *Moraxella catarrhalis*

<400> SEQUENCE: 4

aatgtcaaat cagtcattaa caaagaacaa gtaaatgatg ccaataaaaa gcaaggcatc 60

What we claim is:

1. An immunogenic composition, comprising at least one active component selected from the group consisting of:

(A) an isolated and purified outer membrane protein of a strain of *Moraxella catarrhalis* having a molecular mass of about 200 kDa, as determined by SDS-PAGE; and

(B) a peptide consisting of the amino acid sequence NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys-x-Gln-Gly-Ile (SEQ ID No: 2) or NH₂-Asn-Val-Lys-Ser-Val-Ile-Asn-Lys-Glu-Gln-Val-Asn-Asp-Ala-Asn-Lys (SEQ ID No: 3); and a pharmaceutically acceptable carrier therefor, said at least one active component producing an immune response when administered to a host.

2. The immunogenic composition of claim 1 formulated as a microparticle, capsule, ISCOM, or liposome preparation.

3. The immunogenic composition of claim 1 in combination with a targeting molecule for delivery to specific cells of the immune system or to mucosal surfaces.

4. The immunogenic composition of claim 1 further comprising at least one other immunogenic or immunostimulating material.

5. The immunogenic composition of claim 4 wherein the at least one other immunostimulating material is at least one adjuvant.

6. The immunogenic composition of claim 5 wherein the at least one adjuvant is selected from the group consisting of

40 aluminum phosphate, aluminum hydroxide, QS21, Quil A, calcium phosphate, calcium hydroxide, zinc hydroxide, a glycolipid analog, an octodecyl ester of an amino acid, a muramyl dipeptide and a lipoprotein.

7. The immunogenic composition of claim 6 wherein the host is a primate.

8. The immunogenic composition of claim 7 wherein the primate is a human.

9. A method of generating an immune response in a host, comprising administering thereto an immuno-effective amount of the immunogenic composition of claim 1.

10. The method of claim 9 wherein the immune response is a humoral or a cell-mediated immune response.

11. A method of producing an isolated and purified outer membrane protein of a strain of *Moraxella catarrhalis* having a molecular mass of about 200 kDa, as determined by SDS-PAGE, comprising the steps of:

(a) providing a cell mass of the *Moraxella catarrhalis* strain;

(b) disrupting the cell mass to provide a cell lysate;

(c) fractionating the cell lysate to provide a fraction containing the outer membrane protein substantially free from other cell lysate components, and

(d) recovering said outer membrane protein.

12. The method of claim 11 wherein said *Moraxella catarrhalis* strain is *Moraxella catarrhalis* strain 4223.

13. The method of claim 12 wherein the cell lysate is fractionated by gel electrophoresis.

* * * * *